

Blood fat levels predict diabetes remission following gastric bypass surgery

June 15 2022



Credit: Pixabay/CC0 Public Domain

A newly developed test could help doctors more accurately predict whether individuals who undergo gastric bypass surgery will experience a remission of type 2 diabetes within two years of the procedure,

according to University of Utah Health scientists.

The test, which measures blood levels of ceramides, a waxy, fat-like substance similar to cholesterol, can also be used to assess the likelihood that a [gastric bypass](#) patient will have a recurrence of type 2 diabetes within a decade.

"These [human studies](#) validate our prior findings in rodents that ceramides are major drivers of diabetes. They also reveal a new tool in the physician's arsenal for managing the health of the increasing number of people undergoing [bariatric surgery](#)," says Scott Summers, Ph.D., corresponding author of the study and co-director of the Diabetes Metabolism Research Center.

The study appears in *Med*, a Cell Press [medical journal](#).

In previous studies, Summers and his colleagues found that ceramides can reduce the amount of insulin a body can produce or use, increasing the likelihood that a person will develop type 2 diabetes. Intrigued, they decided to examine the association between ceramides, diabetes, and severe obesity.

More than 24 million Americans are severely obese with a [body mass index](#) (BMI) of 35 or higher, according to the American Society for Metabolic and Bariatric Surgery (ASMBS). As a result, many of them have a substantial risk of developing type 2 diabetes, [heart disease](#), and other severe conditions. For some of these individuals, [gastric bypass surgery](#) is often the best option.

About 250,000 gastric bypass surgeries are performed in the U.S. annually, according to ASMBS. The procedure creates a small pouch from the stomach and connects it directly to the small intestine. As a result, patients will feel full after eating less food.

In some cases, patients can shed more than 30% of their [body weight](#) after surgery and keep it off for at least seven years, according to the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK).

Diabetes is less common after gastric bypass but predicting which patients will have a remission of the disease or a later recurrence has been challenging, says Annelise M. Poss, Ph.D., lead author of the study and a former graduate student in Summers' lab.

To address this challenge, Summers, Poss, and colleagues analyzed blood samples from 1,156 individuals, ages 18-72, collected over a 12-year span. The subjects were divided into three groups: 418 people who underwent gastric bypass surgery; 417 who did not have surgery because they were denied [insurance coverage](#); and 321 severely obese individuals who didn't consider the surgery.

The researchers chose to have two control groups to account for the possibility that those who were denied insurance coverage were more motivated to diet, exercise, and engage in other weight-loss strategies than those who had no interest in gastric bypass

Overall, the scientists found that low [ceramide](#) levels prior to gastric bypass surgery predicted which patients would achieve remission of the diabetes two years later. Of the 67 gastric bypass patients who had diabetes prior to surgery, 49 (73%) either had transient or sustained remission of their type 2 diabetes. However, those individuals with high ceramides did not achieve diabetes remission, regardless of weight change.

"To our knowledge, this is the first study to evaluate the role that ceramides play in type 2 diabetes disease progression in a severely obese population," Poss says. "By measuring ceramides in this population, we

believe we have detected an important biomarker that can be reliably used to better predict the course of type 2 diabetes among patients who undergo gastric bypass."

Among its limitations, the study did not include a group of lean individuals. As a result, they were unable to determine if ceramide levels influence diabetes progression beyond a severely obese population. The researchers also did not prevent participants from pursuing weight loss interventions or lifestyle changes beyond the scope of the study, which focused specifically on Roux-en-Y, the most common gastric bypass [surgery](#).

While the data linking high ceramide levels to diabetes is strong, more research is needed before routine ceramide testing can be recommended, Summers says. As part of that effort, the researchers plan to analyze the association between ceramides levels and diabetes progression in other populations.

"The exciting question we want to answer is this: What are the unique factors that are driving ceramides up in some people who are susceptible to [diabetes](#)?" Summers says. "Clearly, it's not just obesity. A major advance would be to find a family or a unique set of individuals who are prone to accumulate ceramides and develop the disease. That would be a big stride toward developing personalized medical care based on these ceramide biomarkers."

In addition to Summers and Poss, University of Utah Health researchers B. Krick, A. Maschek, B. Haaland, J. Cox, P. Karra, A.R. Ibele, S.C. Hunt, T.D. Adams, W.L. Holland, and M. Playdon contributed to this study. Scientists at the Huntsman Cancer Institute at the University of Utah, Intermountain Healthcare, and Weil Cornell Medicine in Doha, Qatar, also were involved in this research.

More information: Following Roux-en-Y Gastric Bypass Surgery, High Serum Ceramides Demarcate Patients that will Fail to Achieve Sustained Normoglycemia and Diabetes Remission, *Med* (2022). DOI: 10.1016/j.medj.2022.05.011 , [linkinghub.elsevier.com/retrie ...
ii/S2666634022002276](https://linkinghub.elsevier.com/retrieve/pii/S2666634022002276)

Provided by University of Utah Health Sciences

Citation: Blood fat levels predict diabetes remission following gastric bypass surgery (2022, June 15) retrieved 29 June 2024 from <https://medicalxpress.com/news/2022-06-blood-fat-diabetes-remission-gastric.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.