

# Catching COVID-19 in early pregnancy could increase risk of miscarriage

June 21 2022

---



Credit: Pixabay/CC0 Public Domain

Pregnant women who reported being infected with COVID-19 in their first trimester were more likely to have an early miscarriage, finds a new study led by University College London (UCL) researchers.

However, despite the correlation observed in the COVID-19 Contraception and Pregnancy Study (CAP-COVID), researchers say further studies are needed to evaluate any causal association between SARS-CoV-2 infection in early [pregnancy](#) and [miscarriage](#) risk.

The CAP-COVID study, published in *Human Reproduction*, asked 3,041 women who were less than 13 weeks pregnant to record their current and past pregnancy complications, their medical history and whether anyone in their household had been diagnosed, or had symptoms of COVID-19, between 21 May and 31 December, 2020.

The results found that the rate of miscarriage among women who were presumed to be infected (had received a positive COVID-19 diagnosis) was 14%, compared to 8% in those presumed to be uninfected and 5% in those who were uncertain of whether they had the virus.

After adjusting for age, BMI, ethnicity, smoking status, gestational age and the number of previous miscarriages, the risk of early miscarriage appears to be higher (1.7 times more likely) in the "presumed infected" group.

Co-lead author Honorary Associate Professor Dimitrios Mavrelou, who led the project alongside Honorary Senior Lecturer Ephra Yasmin (both UCL EGA Institute for Women's Health & UCLH Reproductive Medicine Unit), said, "As the COVID-19 epidemic reached pandemic status in mid-2020, little was known about the impact of infection on first trimester pregnancy.

"In order to help answer this question we set up an online survey that ran from late May 2020 to December 2020, asking women in early pregnancy to report whether they had suffered a COVID-19 infection and what the outcome of their pregnancy was.

"Over 3,000 women in the first trimester of their pregnancy registered and provided outcomes. Analysis of our data showed a potential association between COVID-19 infection and miscarriage of pregnancy in the first trimester. This is an important public health finding, suggesting that women in early pregnancy may need to be more vigilant with infection prevention measures."

Previous studies have shown that [viral infections](#) during pregnancy such as herpes, HIV-1 and influenza can lead to fetal malformation, preterm labor, growth restriction, stillbirth and spontaneous abortion.

However, this study is the first to investigate the rates of early pregnancy loss during the COVID-19 outbreak to date.

Explaining the potential association between COVID-19 infection and early miscarriage, Honorary Associate Professor Mavrelou said, "Coronaviruses have also been shown to induce a pro-inflammatory cytokine storm in the body. Cytokines are thought to play a role in the development of recurrent pregnancy loss, which could explain the increased risk of early miscarriage in women infected with COVID-19.

"So, although the sample size of women with presumed COVID-19 in this study was too small to reach statistical significance, the findings still highlight the need for pregnant women to exercise social distancing measures and good hygiene throughout their pregnancy to limit their risk of infection."

Researchers relied on self-reported data on early pregnancy loss and SARS-CoV-2 infection without any means of checking validity. Some women in the "presumed uninfected" and "uncertain" groups may have had asymptomatic infections. The number of "presumed infected" in our study was low and therefore the study was relatively underpowered.

**More information:** Neerujah Balachandren et al, SARS-CoV-2 infection in the first trimester and the risk of early miscarriage: a UK population-based prospective cohort study of 3041 pregnancies conceived during the pandemic, *Human Reproduction* (2022). [DOI: 10.1093/humrep/deac062](https://doi.org/10.1093/humrep/deac062)

Provided by University College London

Citation: Catching COVID-19 in early pregnancy could increase risk of miscarriage (2022, June 21) retrieved 24 April 2024 from <https://medicalxpress.com/news/2022-06-covid-early-pregnancy-miscarriage.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.