

A CT algorithm for clear-cell renal cell carcinoma in small solid masses

June 29 2022



Axial CT images in the (A) unenhanced, (B) corticomedullary, and (C) nephrographic phases at the same level depict the mass (arrow, A and C), which shows avid heterogeneous enhancement. Circular ROIs were placed in the most enhancing component of the mass and in ipsilateral renal cortex (circles, B), to determine the mass-to-cortex corticomedullary attenuation ratio. Heterogeneity scores were assigned subjectively using a 5-point Likert type-scale. These two features were used to inform development of the CT score. In this patient, the mass showed avid enhancement (mass-to-cortex corticomedullary enhancement ratio >0.75) and was considered completely heterogeneous (heterogeneous score, 5) by both radiologists, resulting in CT score 5. Nephrectomy revealed clear-cell renal cell carcinoma. Credit: *American Journal of Roentgenology*

According to ARRS' American Journal of Roentgenology (AJR), a 5-tiered CT scoring algorithm may represent a clinically useful tool for diagnosis of clear-cell renal cell carcinoma (RCC) in small (\leq 4 cm) solid renal masses.



"A 5-tiered renal CT <u>algorithm</u>, including mass-to-<u>cortex</u> corticomedullary attenuation ratio and heterogeneity score, had substantial inter-observer agreement, moderate AUC and PPV, and high NPV for diagnosing clear-cell RCC," concluded Nicola Schieda from the department of medical imaging at Canada's Ottawa Hospital.

Schieda and colleagues' study included 148 patients (mean age, 58 years; 73 men, 75 women) with 148 small (\leq 4 cm) solid (>25% enhancing tissue) renal masses that underwent renal-mass CT (unenhanced, corticomedullary, and nephrographic phases) before resection between January 2016 and December 2019. Two <u>radiologists</u> independently evaluated CT examinations and recorded calcification, mass attenuation in all phases, mass-to-cortex corticomedullary attenuation ratio, and heterogeneity score (5-point Likert scale, assessed in corticomedullary phase).

Ultimately, Schieda et al's 5-tiered CT scoring algorithm—including mass-to-cortex corticomedullary attenuation ratio and heterogeneity score—had substantial interobserver agreement (weighted kappa=0.71) and achieved AUC for diagnosing clear-cell RCC of 0.75 (95% CI, 0.68-0.82) for reader 1 and 0.72 (95% CI, 0.66-0.82) for reader 2.

"If validated," the authors of this *AJR* article acknowledged, "the CT algorithm may represent a useful clinical tool for diagnosing clear-cell renal cell carcinoma."

More information: Khalid Al Nasibi et al, Development of a Multiparametric Renal CT Algorithm for Diagnosis of Clear-Cell Renal Cell Carcinoma Among Small (≤4 cm) Solid Renal Masses, *American Journal of Roentgenology* (2022). DOI: 10.2214/AJR.22.27971



Provided by American Roentgen Ray Society

Citation: A CT algorithm for clear-cell renal cell carcinoma in small solid masses (2022, June 29) retrieved 8 May 2024 from https://medicalxpress.com/news/2022-06-ct-algorithm-clear-cell-renal-cell.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.