


Cutting-edge imaging yields new insights into stroke

June 9 2022, by Greg Basky



Hemorrhage biomarkers provide insights into the extent of heme biotransformation during recovery as well as the extent of hemorrhage-induced oxidative damage. Credit: *Metallomics* (2022). DOI: 10.1093/mtomcs/mfac007

Hemorrhagic stroke, where a weakened vessel in the brain ruptures, can lead to permanent disability or death. Across the globe, over 15 million people are coping with its effects.

A study by researchers from the University of Saskatchewan (USask) and Curtin University in Australia has moved us one step closer to identifying when the bleeding associated with a hemorrhagic stroke starts—critical information for improving patient outcomes.

Time is of the essence when it comes to stroke; the sooner doctors can start treatment, the better the odds they can limit damage.

Using the Mid-IR beamline at the Canadian Light Source at USask, the team examined brain tissue samples with a special technique called Fourier-transform infrared imaging. The researchers were led by Dr. Lissa Peeling, a neurosurgeon at the Royal University Hospital and an Associate Professor in the Department of Surgery at USask.

The novel approach enabled the researchers to identify changes in the brain specific to hemorrhagic stroke.

Dr. Jake Pushie, a member of Dr. Kelly's and Dr. Peeling's research team at USask's College of Medicine, said the combination of the beamline and infrared imaging made it easy to detect markers of [brain damage](#) caused by hemorrhagic stroke.

"In a sense, this is giving us 'superhuman vision' to look at these brains and map out what's happening metabolically," said Pushie.

With synchrotron technology, the team could see where a bleed originated and the extent of oxidative damage it caused—something impossible to do with a microscope or traditional approaches to imaging. Their findings were published in *Metallomics*.

Armed with this new approach, and a better understanding of what they are looking for, Pushie and colleagues will now go back through their extensive "library" of stroke tissue samples to gain a clearer picture of

the speed at which [oxidative damage](#) begins to ramp up.

The team's findings could eventually enable doctors to use clinical imaging—such as MRI or CT scans—to pinpoint where, and how long ago, a [hemorrhagic stroke](#) occurred in the [brain](#).

Miranda Messmer is another member of the team who was a USask undergraduate research student with the College of Arts and Science at the time of the study and will be a graduate student with the USask College of Medicine in the fall.

Messmer said knowing when bleeding has started can provide clinicians with a clearer picture of the time window they have to act.

"Being able to understand what is going on biologically, when we see any kinds of changes in the clinical images, could help doctors provide better care when it comes to minimizing the tissue damage associated with stroke," said Messmer.

More information: M J Pushie et al, Multimodal imaging of hemorrhagic transformation biomarkers in an ischemic stroke model., *Metallomics* (2022). [DOI: 10.1093/mtomcs/mfac007](https://doi.org/10.1093/mtomcs/mfac007)

Provided by Canadian Light Source

Citation: Cutting-edge imaging yields new insights into stroke (2022, June 9) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-06-cutting-edge-imaging-yields-insights.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.