

Understanding C. diff infection

June 6 2022, by Laurel Kelly



A medical illustration of *Clostridioides difficile* bacteria, formerly known as *Clostridium difficile*, presented in the Centers for Disease Control and Prevention (CDC) publication entitled, Antibiotic Resistance Threats in the United States, 2019. Credit: CDC

Clostridioides difficile is a bacterium that causes an infection of the large intestine. The bacterium is often referred to as *C. difficile* or *C. diff*. Some people carry *C. diff* bacteria in their intestines but never become sick. They are carriers of the bacteria and may spread infections.

Illness from *C. diff* typically occurs after using antibiotics. It most commonly affects [older adults](#) in hospitals or long-term care facilities. About 200,000 people in the U.S. are infected annually with *C. diff* in a hospital or long-term care setting. These numbers are lower than in previous years because of improved prevention measures.

Signs and symptoms of *C. diff* infection usually develop within five to 10 days after starting a course of antibiotics. However, they may occur as soon as the first day or up to three months later.

The most common signs and symptoms of mild to moderate *C. diff* infection are watery diarrhea and mild abdominal cramps. Additional signs and symptoms of severe infection include rapid heart rate, dehydration, fever and nausea.

Complications of *C. diff* infection include dehydration, [kidney failure](#), colon damage and death.

A person is treated only when showing signs or symptoms of infection. Treatments can include antibiotics and surgery to remove the diseased portion of the colon. People who carry the bacteria but are not sick are not treated.

Approximately 25% of people treated for *C. diff* infection get sick again, either because the [initial infection](#) never went away or because they've been reinfected with a different strain of the bacteria. The risk increases with each *C. diff* infection episode, with the risk of getting

sick again exceeding 50% after three or more infections. Additional treatment options for recurrent disease may include fecal microbiota transplant.

Provided by Mayo Clinic

Citation: Understanding C. diff infection (2022, June 6) retrieved 6 August 2024 from <https://medicalxpress.com/news/2022-06-diff-infection.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--