


Primary hyperparathyroidism linked to bone fracture, stroke and heart attack

June 16 2022



Cumulative incidences for patients with primary hyperparathyroidism (pHPT) and control individuals. The cumulative incidence of events for untreated pHPT and control groups was estimated using 1-minus Kaplan-Meier estimate of the corresponding survival function and presented with 95% CIs (indicated by shaded areas). Credit: *JAMA Network Open* (2022). DOI: 10.1001/jamanetworkopen.2022.15396

In the hormonal disorder primary hyperparathyroidism, patients' risk of hip fracture is elevated by 51%, and that of heart attack or stroke 45%. These are results from a major study from the University of Gothenburg, Sweden, which also demonstrates that surgery is associated with reduced risks of these outcomes.

Primary hyperparathyroidism is a common hormone disorder, especially among the elderly. It occurs more often in women, and some 3% of [postmenopausal women](#) are affected. Rather than causing specific symptoms, it is often detected by chance in connection with blood tests, which then show elevated [calcium levels](#) and normal or elevated parathyroid hormone levels. This disturbed calcium balance in the blood can cause kidney, skeletal, and cardiovascular damage.

Previous studies have linked primary hyperparathyroidism to osteoporosis and [cardiovascular disease](#). However, since these studies have been few and small, the association has been debated.

Many complications

This study, published in the journal *JAMA Network Open*, is based on national register data from the Swedish National Board of Health and Welfare. All the 16,374 patients included were diagnosed with primary hyperparathyroidism sometime between 2006 and 2017. Each was compared with ten control individuals from the population born in the same year, of the same sex, and residing in the same county.

"We show that untreated primary hyperparathyroidism means a 51% higher risk of hip fracture and a 45% increase in [heart attack](#) or stroke risk. The kidney stone risk is almost quadrupled and, additionally, the risk of death is raised by 72%. The increased likelihood of these

complications highlights the importance of identifying patients with this hormonal disease," says Kristian Axelsson, researcher at the University of Gothenburg, resident in general medicine within the public primary care in Region Västra Götaland, and the study's first author.

Surgery reduces risks

An operation to remove the [parathyroid glands](#) (parathyroidectomy), entirely or partially, is the only definitive treatment for primary hyperparathyroidism. Patients who meet special criteria may undergo [surgery](#). Careful monitoring is recommended for those who do not.

The study shows that the elevated risks of [hip fracture](#), heart attack, and stroke decreased significantly for patients who underwent surgery.

"The study indicates that surgery clearly reduces the risk of osteoporosis fractures, fall injuries, and death in cardiovascular events, and these are vital findings that may lead to more patients being selected for surgery. Our results support the view that patients' cardiovascular risk, as well as the high fracture risk, should be considered in surgery decisions," says Mattias Lorentzon, professor of geriatric medicine at the University of Gothenburg, chief physician at Sahlgrenska University Hospital, and senior author of the study.

More information: Kristian F. Axelsson et al, Analysis of Comorbidities, Clinical Outcomes, and Parathyroidectomy in Adults With Primary Hyperparathyroidism, *JAMA Network Open* (2022). [DOI: 10.1001/jamanetworkopen.2022.15396](https://doi.org/10.1001/jamanetworkopen.2022.15396)

Provided by University of Gothenburg

Citation: Primary hyperparathyroidism linked to bone fracture, stroke and heart attack (2022, June 16) retrieved 12 May 2024 from <https://medicalxpress.com/news/2022-06-primary-hyperparathyroidism-linked-bone-fracture.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.