

Study reveals deep inequalities of care for atrial fibrillation patients

June 7 2022, by Mike Addelman



Credit: Pixabay/CC0 Public Domain

Many ethnic minority patients in England with the most common form of atrial fibrillation (AF) are not being prescribed the blood thinners they need to lower their stroke risk, according to data scientists at the

University of Manchester.

The study compared recommended treatment (oral anticoagulation such as warfarin), less effective treatment (aspirin), and no treatment.

Compared to white patients, Black and other ethnic minorities patients were 22%, and 24% less likely to receive oral anticoagulation, respectively. These estimates have taken into account patient age, sex, comorbidities, residence deprivation (ranked 1 to 5, from least to most deprived), and bleeding [risk factors](#).

The study of the heart condition, published in *PLOS Medicine* today (June 7, 2022), also identifies people from deprived neighborhoods as being less likely to receive the treatment.

Patients living in the most deprived areas were 15% less likely to receive oral anticoagulation, compared to patients living in the least deprived areas (on top of the ethnicity disparities).

For example, a black patient living in deprivation quintile 5 was 16% likely to be prescribed aspirin-only and 62% [oral anticoagulants](#), while respective estimates for a white patient living in deprivation quintile 1 were 12% and 72%.

Adjusting for patient characteristics, practices in London, the North West, and Yorkshire and Humber generally performed worse than other regions, with a no-treatment probability of around 18% compared to practices in the North East and the South West regions with a no-treatment probability of around 14%.

And the team also identified an association between different comorbidities such as dementia, liver disease and cancer and under-prescribing.

Patients with dementia, [liver disease](#), malignancy, and history of falls were associated a 48%, 42%, 26% and 18% lower prescribing rate of [oral anticoagulation](#), respectively.

Patients with history of ischemic heart disease were also more likely to receive aspirin-only, even though this is not considered best practice for stroke prevention.

The data also showed that incidence rates of [atrial fibrillation](#) rose by almost a quarter in England between 2009 and 2019, and were markedly higher in men than in women.

The researchers analyzed almost 200,000 patients aged 18+ and registered with an English general practice between 2009 and 2019, contributing data to the U.K. Clinical Practice Research Datalink database.

However, they also showed there was an overall rise in the proportion of patients with AF prescribed oral anticoagulants increased substantially between 2011 and 2015.

But the increase, they said, may be related to a change in European Society of Cardiology guidelines in 2010, which recommended anticoagulation for all patients with AF at moderate-risk to high-risk of stroke.

That was followed by a 2012 update recommending avoidance of aspirin prescribing in low-stroke risk patients.

Lead author Alyaa Ajabnoor, a Ph.D. researcher from The University of Manchester said: "Atrial fibrillation is associated with increased risk of stroke, cardiovascular comorbidities, and mortality, and currently accounts for 1% of the total healthcare expenditure in the United

Kingdom.

"The treatment of choice is anticoagulation, however many patients with non-valvular atrial fibrillation -the most common form of the disease—are not receiving the therapy they need.

"Despite an increase in prescribing, we report important racial and socioeconomic inequalities in the prescribing of oral anticoagulants.

"People with [low socioeconomic status](#) and black or other non-white ethnicities were associated with the prescription of aspirin-only or no-treatment at all compared to [white patients](#) or those with higher socioeconomic status."

Co-author Professor Evan Kontopantelis from The University of Manchester said: "Though this study depends on accurate recording of conditions by [health professionals](#), we feel the overriding findings are robust and stark.

"Our findings suggest that to improve atrial fibrillation outcomes, these inequalities need to be addressed through equitable interventions to improve anticoagulant prescribing to prevent strokes and reduce mortality."

More information: Incidence of non-valvular atrial fibrillation and oral anti-coagulant prescribing in England, 2009 to 2019: a cohort study, *PLOS Medicine*

Provided by University of Manchester

Citation: Study reveals deep inequalities of care for atrial fibrillation patients (2022, June 7)

retrieved 19 April 2024 from

<https://medicalxpress.com/news/2022-06-reveals-deep-inequalities-atrial-fibrillation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.