


Suicidal thoughts, behaviors linked to hormone-sensitive brain disorder

June 13 2022



Prevalence of self-injurious thoughts and behaviors in those with prospectively confirmed PMDD, subdivided by psychiatry comorbidity status. Credit: Tory Eisenlohr-Moul et al, *BMC Psychiatry* (2022). DOI: 10.1186/s12888-022-03851-0

A new global study published in *BMC Psychiatry* reports that 34% of people with premenstrual dysphoric disorder have attempted suicide.

The study is the largest of its kind to examine rates of [suicidal thoughts](#) and behaviors among people diagnosed with the disorder by a [health care provider](#) based on daily symptom ratings, the only reliable method currently available.

Because previous studies of suicide and ideation have relied on less valid self-reported measures of [premenstrual dysphoric disorder](#), which is commonly called PMDD, the new findings offer the strongest scientific evidence to date that the disorder is likely an independent contributor to suicidal thoughts and actions.

"We've uncovered an extremely worrying rate of suicide ideation and attempts among those with PMDD, highlighting the need to take this issue seriously," said Tory Eisenlohr-Moul, assistant professor of psychiatry at the University of Illinois Chicago and lead author of the study. "These findings offer powerful evidence that the link between PMDD and suicide is independent of depression, post-[traumatic stress disorder](#), or other [mental health conditions](#) that are known to increase ideation and attempts."

Premenstrual dysphoric disorder is a cyclical, hormone-based disorder that impacts approximately 1 in 20 reproductive-age women who were assigned female at birth. The condition is often underdiagnosed, misdiagnosed or dismissed entirely by [medical professionals](#), despite patient reports of debilitating anxiety, hopelessness and a variety of physical symptoms in the two weeks before menses.

To better understand PMDD, the researchers analyzed information from the Global Survey of Premenstrual Disorders, which included 3,153 people from over 56 countries and more than 2,000 responses. The survey was conducted by the International Association for Premenstrual Disorders, Me v PMDD and Vicious Cycle to help understand the scope of premenstrual disorders and the impact of PMDD.

Analysis of the information provided by the 599 respondents who reported a prior daily ratings-based diagnosis of PMDD by a health care provider (23% of respondents) revealed that 34% have attempted suicide during a PMDD episode. On average, patients waited 12 years and saw around six providers before receiving an accurate diagnosis of PMDD.

The data also showed high rates of lifetime active suicidal ideation (72%), planning (49%), intent (42%) and preparing (40%) for an attempt, and non-suicidal self-injury (51%) among patients with PMDD diagnoses. Low-to-moderate income, history of major depression or post-traumatic stress disorder and nulliparity—never giving birth—were predictors of lifetime active suicidal ideation and attempts. Older age and borderline personality disorder were additional predictors of lifetime attempts.

Rates of self-injurious thoughts and behaviors also were broken down by those with only PMDD and those with PMDD who also reported having received at least one other mental health diagnosis, like depression. Even among those who had never received another mental health diagnosis, rates were high: 67% reported active suicidal ideation, compared to 74% who also had psychological comorbidity, for example.

Eisenlohr-Moul said she'd expect to see a far greater difference in rates between categories if thoughts and behaviors were only, or even mostly, due to other underlying mental health problems. The data suggests to her that women who are neurobiologically sensitive to hormone changes, as in the case with PMDD, may be at increased risk for suicidal thoughts and behavior.

"One of the big challenges with PMDD is that the [medical community](#) has not just been slow to understand this condition but even to believe it exists," said Eisenlohr-Moul, who is also chair of IAPMD's clinical advisory board. "Providers and communities often dismiss patients'

concerns, in part because women's complaints are less likely to be taken seriously than men's but also because of persistent and even sexist stigma and misconceptions around menstruation in general."

"PMDD is not a hormone imbalance. It is a neurobiological sensitivity to natural and normal changes in progesterone and estrogen levels," she said.

"Our study reveals just how destructive PMDD is," said Sandi MacDonald, co-founder and executive director of the International Association for Premenstrual Disorders. "This is a galvanizing movement in women's health. PMDD is a perfect storm where #MeToo and #TimesUp, meet mental health awareness, meets suicide prevention."

While PMDD has been included in the Diagnostic and Statistical Manual of Mental Disorders since 2013 as a [major depressive disorder](#), there is still no recommended standard screening of suicidal ideation in patients with the condition.

More information: Tory Eisenlohr-Moul et al, Prevalence of lifetime self-injurious thoughts and behaviors in a global sample of 599 patients reporting prospectively confirmed diagnosis with premenstrual dysphoric disorder, *BMC Psychiatry* (2022). [DOI: 10.1186/s12888-022-03851-0](#)

Provided by University of Illinois at Chicago

Citation: Suicidal thoughts, behaviors linked to hormone-sensitive brain disorder (2022, June 13) retrieved 24 May 2024 from <https://medicalxpress.com/news/2022-06-suicidal-thoughts-behaviors-linked-hormone-sensitive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.