


New technique for detecting typhoid infections faster, which is more accurate than conventional testing

June 30 2022



Estimated seroincidence of typhoidal *Salmonella* by study community and age group. Age groups are denoted by point shapes for the median, with lines indicating the 95% CI. Boxes reflect the height of the median estimate for the overall population-based serosurvey. Credit: *The Lancet Microbe* (2022). DOI: 10.1016/S2666-5247(22)00114-8

An international team of researchers found a new technique for detecting typhoidal Salmonella infections that can significantly help vaccination planning in countries where typhoid is prevalent. The new approach detects typhoid antibodies in dried blood spots. It is faster and more accurate than blood cultures and simplifies disease detection in remote areas.

The collaborative study includes authors from Bangladesh, Nepal, Pakistan, Ghana, Canada, South Korea, England and Germany. It was published in *The Lancet Microbe*.

"Typhoid is difficult to diagnose, and we know the burden is much higher than what we find through routine hospital surveillance," said Kristen Aiemjoy, first author of the study. Aiemjoy is an assistant professor in the Division of Epidemiology, Department of Public Health Sciences at UC Davis. "You can't diagnose it on symptoms. The only available test is expensive, slow, not very widely available and has only around 60% sensitivity."

The World Health Organization (WHO) estimates that 11 to 20 million people get sick from [typhoid](#), and between 128,000 and 161,000 people die from it every year. The life-threatening infection is caused by the bacterium Salmonella Typhi and is usually spread through contaminated food or water. It is most common in parts of the world with poor sanitation.

The WHO recommends typhoid vaccines to control endemic and epidemic typhoid fever.

Many countries lack the infrastructure to conduct comprehensive typhoid testing, particularly in rural areas. As a result, the incidence in Bangladesh, Nepal, Pakistan and other countries may be significantly undercounted, making it difficult to develop effective vaccination

strategies.

Dried blood samples reveal typhoid infections

The team developed a tool that relies on antibody responses from dried capillary [blood samples](#). The blood is collected using a finger prick, similar to the small amount used for a blood glucose monitor. This assay is easy to perform, can be readily accomplished in remote, rural areas and is far more accurate than blood cultures.

To develop this technology, the team collected [blood spots](#) from 1,420 people with confirmed typhoid infections in Bangladesh, Ghana, Nepal and Pakistan. They tested the samples for antibody responses against two antigens: Hemolysin E (HlyE) and Salmonella lipopolysaccharide (LPS). The researchers showed the antibody responses to both HlyE and LPS persisted for months after infection, making these antibodies excellent biomarkers to measure typhoid incidence.

In addition, the researchers developed algorithms to determine when a person contracted typhoid, allowing them to measure the disease's incidence in populations over time. To accomplish this, they measured [antibody responses](#) in a population-based sample of 1,740 children and young adults in Nepal, Pakistan, Bangladesh and Ghana. When comparing their results to existing blood culture data, they found that the true number of typhoid infections was much higher across countries and regions.

"Instead of just taking a sample, looking at the person's antibody response and saying they're negative or positive, we can model the antibody decay," said Aiemjoy. "We use that decay rate to infer when those people were most likely exposed, which provides vital information about the force of infection."

New tool allows broader surveillance of disease

Current typhoid estimates generally come from capital cities, often the only areas with blood culturing capabilities. This leaves large regions where public health officials can only guess disease prevalence. In addition to providing more comprehensive information outside cities, the new method can also pinpoint subpopulations with high percentages of typhoid in urban areas.

"The approach described in this study has the potential to expand the geographical scope of typhoidal Salmonella surveillance and generate incidence estimates that are comparable across geographical regions and time," said Farah Naz Qamar, Jessani Endowed Professor at Aga Khan University in Pakistan and one of the senior authors on the paper. "This will have a great impact on public health, as it can guide the possible introduction of typhoid conjugate vaccines in national programs for typhoid amelioration where needed the most."

Many well-known infectious disease researchers contributed to the study, including Samir Saha at the Child Health Research Foundation in Bangladesh, Jason Andrews and Steve Luby at Stanford University and Richelle Charles at Mass General Hospital. Denise Garrett, at the Sabin Vaccine Institute, was the principal investigator.

While the study is limited to typhoid, the authors believe this approach could be applied to other [infectious diseases](#), revolutionizing how nations conduct epidemiological surveillance.

"This is a big jump for typhoid surveillance, but it's also a big methodological jump in seroepidemiology," said Aiemjoy. "We are now planning on applying this to other infectious diseases like scrub typhus and melioidosis."

More information: Kristen Aiemjoy et al, Estimating typhoid incidence from community-based serosurveys: a multicohort study, *The Lancet Microbe* (2022). [DOI: 10.1016/S2666-5247\(22\)00114-8](https://doi.org/10.1016/S2666-5247(22)00114-8)

Provided by UC Davis

Citation: New technique for detecting typhoid infections faster, which is more accurate than conventional testing (2022, June 30) retrieved 18 April 2024 from <https://medicalxpress.com/news/2022-06-technique-typhoid-infections-faster-accurate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.