


# Adolescent childbirth remains linked to poor outcomes for both mother and child in Cote d'Ivoire

July 6 2022



Flow chart of patients. Credit: *Gynecology and Obstetrics Clinical Medicine* (2022). DOI: 10.1016/j.gocm.2022.04.002

According to the 2014 general population census for Cote d'Ivoire, one in four of the country's women aged 20 to 24 years gave birth to at least one child before they reached 18 years. A retrospective study of teenage childbirth in the west African country has found that these adolescent

births are often associated with a range of risks for both mother and child.

Researchers from the Department of Obstetrics and Gynecology at the Cote d'Ivoire's University of Abidjan-Yopougon, looked at three years' worth of data from their maternity hospital. They compared the experiences of an adolescent group of 1,040 girls aged 10 to 19 years, with a 736-strong young adult group aged 20 to 24 years. Their results have now been published in the KeAi journal *Gynecology and Obstetrics Clinical Medicine*.

The average age of the [adolescent girls](#) they studied was 17.45 years, with the youngest aged 12 years. They found that this group accessed help from a consultant less often (and later) than the young adult group. And the proportion of adolescent girls (47.6%) versus young adults (39.1%) who underwent a cesarean section was significantly higher.

According to study corresponding author EdeleKacou Aka: "We also found that the adolescents were 3.5 times more likely to develop [high blood pressure](#) and its complications in the weeks following [birth](#), and 2.5 times more likely to contract an infection during this (postpartum) period."

In addition, the risks of premature birth, the baby dying within 28 days of birth, and the mother or child being admitted to hospital following the birth, were all significantly higher in the adolescent group. The adolescent girls were also more likely to have limited or narrowed pelvic bone.

Aka adds: "These data show that adolescent childbirth remains associated with a poor maternal-fetal prognosis. And they highlight the need for more education on the [prenatal care](#) available, along with policies encouraging contraceptive methods to reduce adolescent

pregnancies and childbirth."

**More information:** Edele Kacou Aka et al, Obstetrical specificities of Ivorian adolescent mothers on a single-center hospital series of 1040 cases, *Gynecology and Obstetrics Clinical Medicine* (2022). [DOI: 10.1016/j.gocm.2022.04.002](https://doi.org/10.1016/j.gocm.2022.04.002)

Provided by KeAi Communications Co.

Citation: Adolescent childbirth remains linked to poor outcomes for both mother and child in Cote d'Ivoire (2022, July 6) retrieved 11 July 2024 from <https://medicalxpress.com/news/2022-07-adolescent-childbirth-linked-poor-outcomes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--