


Coinfection with influenza A could suppress replication in SARS-CoV-2

July 12 2022



3D print of influenza virus. The virus surface (yellow) is covered with proteins called hemagglutinin (blue) and neuraminidase (red) that enable the virus to enter and infect human cells. Credit: NIH

Coinfection of SARS-CoV-2 and influenza A virus changes neither the trajectory, nor the severity of influenza A virus, regardless of timing.

But should the host contract influenza A virus first, the response to that infection can significantly suppress SARS-CoV-2, according to research published this week in the *Journal of Virology*, a publication of the American Society for Microbiology.

"The research is important, because the human population now has 2 circulating respiratory RNA viruses with high pandemic potential: SARS-CoV-2 and [influenza A](#)," the investigators wrote. "As both viruses infect the airways, and can result in significant morbidity and mortality, it is imperative that we also understand the consequences of coinfection."

Several [clinical studies](#) had previously reported on co-infection of SARS-CoV-2 with other viruses. "In particular, coinfection with SARS-CoV-2 and influenza A [virus](#) was common early in the COVID-19 pandemic, prior to the enforcement of masks and social distancing," said corresponding author Benjamin R. tenOever, Ph.D., professor of microbiology, New York University, Langone Health, New York, N.Y.. These viruses infect the same cells within the airway.

Notably, the investigators found that influenza A virus interferes with SARS-CoV-2 replication in the lung and can continue to do so even more than 1 week after clearance of influenza A according to the research.

"These data suggest the presence of factors intrinsic to or induced by [influenza A virus] that may restrict the growth of SARS-CoV-2, but it remains unclear whether this effect plays a role on disease severity," the researchers wrote.

The investigators performed the experiments in cultured cells, as well as in a golden hamster animal model. "... animals were administered the 2 viruses simultaneously, and examined at days 1, 3, 5, 7 and 14 post infection," said tenOever. The researchers also conducted experiments in

which they first challenged the animals with either virus, followed three days later by the other virus, monitoring [them] at days 1, 3, and 5 post-second challenge.

"This study could be used as an example of how an [immune response](#) to something unrelated can provide protection against SARS-CoV-2," said tenOever.

The team demonstrated that coinfection does not result in a worse outcome of disease in an animal model. "These results suggest that coinfection with SARS-CoV-2 and influenza A virus does not represent a looming threat for humanity," said tenOever.

More information: Kohei Oishi et al, The Host Response to Influenza A Virus Interferes with SARS-CoV-2 Replication during Coinfection, *Journal of Virology* (2022). [DOI: 10.1128/jvi.00765-22](https://doi.org/10.1128/jvi.00765-22)

Provided by American Society for Microbiology

Citation: Coinfection with influenza A could suppress replication in SARS-CoV-2 (2022, July 12) retrieved 17 May 2024 from <https://medicalxpress.com/news/2022-07-coinfection-influenza-suppress-replication-sars-cov-.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|