


# Difficulty swallowing linked to chronic opioid use

July 28 2022, by Bill Snyder

---



Credit: Pixabay/CC0 Public Domain

Chronic use of opioid painkillers like hydrocodone and oxycodone significantly increases the risk of dysphagia, difficulty swallowing, according to researchers at Vanderbilt University Medical Center.

While opioid use was not associated with rare swallowing disorders such as achalasia, [patients](#) on opioids had a significantly higher burden of symptoms than non-users, and their symptoms tended to be more persistent, Dhyanes Patel, MD, and colleagues reported in the journal

*Gastroenterology.*

The [retrospective study](#) included 4,075 patients who had undergone high-resolution manometry (HRM), of whom 869 were being treated chronically with [opioid drugs](#) for pain. HRM is a diagnostic technique that measures the motility of the esophagus.

Dysphagia was diagnosed in 65% of patients on chronic opioids, compared to 51% of those who were not taking opioid drugs. Hydrocodone was the most-commonly used painkiller in the opioid group, followed by [oxycodone](#), tramadol and morphine.

This was the largest cohort of patients studied to date to determine the impact of chronic opioid use on esophageal function, said Patel, assistant professor of Medicine in the Vanderbilt Center for Swallowing and Esophageal Disorders and the paper's corresponding author.

Tramadol, which does not provide the full analgesic action of other opioids, was not associated with abnormal HRM findings or a hypercontractile esophagus. It could be used to reduce the risk of symptoms, the researchers suggested.

Given the significant symptom burden in this group of patients, a more "definitive" therapeutic approach such as POEM (an endoscopic technique) should be offered to those on [opioids](#) who are diagnosed with achalasia, they added.

With nearly 4% of the U.S. population under chronic opioid treatment for non-cancer-related chronic pain, these findings highlight the importance of educating and counseling patients about how chronic [opioid use](#) might affect their symptoms long term, the researchers concluded.

**More information:** Dhyanesh A. Patel et al, Opioid Exposure Differentially Impacts Esophageal Body Contraction Over the Lower Esophageal Sphincter, *Gastroenterology* (2022). [DOI: 10.1053/j.gastro.2022.04.051](https://doi.org/10.1053/j.gastro.2022.04.051)

Provided by Vanderbilt University

Citation: Difficulty swallowing linked to chronic opioid use (2022, July 28) retrieved 11 May 2024 from <https://medicalxpress.com/news/2022-07-difficulty-swallowing-linked-chronic-opioid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.