

Lower fecundability with residence in disadvantaged neighborhood

July 1 2022



Residence in a socioeconomically disadvantaged neighborhood is

associated with reduced fecundability, according to a study published online June 30 in *JAMA Network Open*.

Mary D. Willis, Ph.D., from Oregon State University in Corvallis, and colleagues examined the association between residence in disadvantaged [neighborhoods](#) and fecundability in a prospective preconception cohort study involving 6,356 participants who identified as female, were age 21 to 45 years of age, were attempting [conception](#) without fertility treatment, and who provided a valid residential address. Fecundability ratios were estimated for the associations between a standardized area deprivation index (ADI), applied to each residential address, and fecundability.

The researchers observed 3,725 pregnancies for 27,427 menstrual cycles of follow-up. The adjusted fecundability ratios were 0.79 and 0.77 for national-level and within-state ADI rankings, respectively, comparing the top and bottom deciles for disadvantaged neighborhood status. Some evidence of nonlinearity was seen in the association with restricted cubic splines. Among participants with lower annual incomes (

Citation: Lower fecundability with residence in disadvantaged neighborhood (2022, July 1) retrieved 6 May 2024 from

<https://medicalxpress.com/news/2022-07-fecundability-residence-disadvantaged-neighborhood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.