


A methodological leap in the exploration of memory

July 27 2022




Mobility of glutamate receptors on the surface of a rat hippocampal neuron, measured by tracking individual molecules. Credit: © Benjamin Compans and Daniel Choquet / IINS / CNRS-Université de Bordeaux

Neurons communicate with each other across synapses, areas of close contact where neurotransmitter molecules released from one neuron act on receptors embedded in the membrane of the opposite neuron. Previous research conducted by the team of Daniel Choquet, researcher at the CNRS and Director of the Interdisciplinary Institute for

Neurosciences (CNRS/University of Bordeaux) has discovered that these receptors are not stationary, but instead move constantly in the membrane. The same scientists suggested and indirectly demonstrated that this movement modifies the number of receptors in a synapse at a given time to modulate the effectiveness of synaptic transmission and, as a result, certain types of learning and memory.

Until now, however, it was not possible to observe receptor mobility in situations more natural than neuron cultures. This has now been achieved thanks to the development of a comprehensive toolbox with which scientists have established that this mobility exists in intact brain tissue, and that it is indispensable for certain types of memory, such as the contextual fear memory tested in this study.

This toolbox consists of a new animal model, improved high-resolution imaging technology and techniques for labeling and controlling receptor dynamics. It will allow the study of any region of the brain in addition to the hippocampus, can be transposed to other types of [receptors](#), and will be used by the team to study the possible role of receptor mobility in [intellectual disabilities](#) and [autism spectrum disorders](#).



A neuron in a hippocampal slice from the mouse model developed by the team, in which the glutamate receptor GluA2 is 'labelled' to create a stain of an isolated neuron. Magenta: stain of GluA2, on the dendrites of the neuron; green: Soluble GFP stain of the cell body and axon. Images of the neuron are produced with a light sheet microscope, a high-resolution imaging technology. Credit: © Angela Getz, Mathieu Ducros, Daniel Choquet / IINS & BIC / CNRS-Université de Bordeaux-Inserm.

More information: Angela M. Getz et al, High-resolution imaging and manipulation of endogenous AMPA receptor surface mobility during

synaptic plasticity and learning, *Science Advances* (2022). DOI: [10.1126/sciadv.abm5298](https://doi.org/10.1126/sciadv.abm5298).
www.science.org/doi/10.1126/sciadv.abm5298

Provided by CNRS

Citation: A methodological leap in the exploration of memory (2022, July 27) retrieved 4 May 2024 from <https://medicalxpress.com/news/2022-07-methodological-exploration-memory.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.