

Monkeypox virus is frequently detected in saliva, semen and other clinical samples from infected patients

July 14 2022



Credit: Pixabay/CC0 Public Domain

Viral DNA can be frequently detected in different clinical samples from

monkeypox-infected patients, including saliva and semen, according to a new study led by the Barcelona Institute for Global Health (ISGlobal), a center supported by the "la Caixa" Foundation, and the Hospital Clínic of Barcelona. The study, published in *Eurosurveillance*, contributes to a better understanding of how this emerging disease is transmitted.

The current monkeypox outbreak represents another zoonotic disease that has crossed borders. In the last six months, more than 9,000 cases of monkeypox have been reported worldwide in countries where the disease is not endemic. Initial cases were detected in Britain, Portugal and Spain, mainly in men who have sex with men. However, the disease has spread to many other countries and there is concern that it may spread to vulnerable populations, such as immunocompromised patients or children.

The disease is known to be transmitted by direct contact with the lesions of an infected patient or by surfaces contaminated by these lesions, but little is known about the possible presence of virus in other biological samples, such as [saliva](#), urine or semen.

In this study, the team led by Mikel Martinez, ISGlobal researcher, and Jose Luis Blanco (Hospital Clinic), investigated the presence of genetic material of the virus in different [biological samples](#), collected at different times, from 12 patients with confirmed monkeypox infection. At the time of diagnosis, a high viral DNA load was detected in the skin lesions of all patients. In addition, DNA was detected in the saliva of all cases, some of them with high viral loads. Only one previous study had tested saliva, in one single patient. Viral DNA was also detected in rectal (11 of 12 patients), nasopharyngeal (10/12 patients), semen (7/9 patients), urine (9/12 patients) and fecal (8/12 patients) samples.

"A couple of previous studies had already shown occasional presence of viral DNA in some samples and in some patients, but here we show that

viral DNA is frequently present in various biological fluids, particularly saliva, during the acute phase of the disease, and up to 16 days after the onset of symptoms in one patient," explains Aida Peiró, first author of the study.

The authors point out that the presence of viral DNA does not necessarily mean infectious virus, and that the next step will be to try to isolate infectious virus from such samples. However, the high viral load detected in saliva or semen suggests that such fluids have infectious potential, they add.

"The results of our study contribute to a better understanding of the mechanisms and dynamics of [virus](#) transmission, as well as the possible role of sexual transmission," Martinez concludes.

More information: Mikel Martinez et al, Frequent detection of monkeypox virus DNA in saliva, semen, and other clinical samples from 12 patients, *Eurosurveillance* (2022). [DOI: 10.2807/1560-7917.ES.2022.27.28.2200503](https://doi.org/10.2807/1560-7917.ES.2022.27.28.2200503)

Provided by Barcelona Institute for Global Health

Citation: Monkeypox virus is frequently detected in saliva, semen and other clinical samples from infected patients (2022, July 14) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-07-monkeypox-virus-frequently-saliva-semen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.