

Neurons specialized in encoding sound emerge before birth

July 11 2022



3D picture of the neurosensory elements of a mouse cochlea soon after birth (neurons are at the centre and hair cells in the periphery, in a spiral shape). Credit: Csaba Ádori



Distinct neuron types in the auditory organ are necessary for encoding different features of sound and relaying them to the brain. Researchers at Karolinska Institutet provide evidence of an early, neuronal activity-independent, emergence of the different subtypes of auditory neurons, prior to birth in mice. The findings have recently been published in *Nature Communications*.

Previous studies have provided ambiguous results on whether the different subtypes of <u>auditory neurons</u> emerge during prenatal or postnatal development, with in the latter case, a possible role of neuronal activity in generating their diversity. In this new study, researchers demonstrate that the fate of auditory neuron subtypes is under genetic control in the prenatal period, and reveal the complex <u>molecular</u> <u>networks</u> controlling their genesis.

Why are these results important?

"Our findings demonstrate that most aspects directly linked to the functional diversity of auditory neurons seem to be intrinsically, molecularly defined before birth," says Francois Lallemend, Associate Professor at the Department of Neuroscience, and last author of the article. "This has implications on how we can interpret peripheral auditory (dys-)functions at birth but also for devising ways to convert neuron types in pathological conditions or congenitally deaf cochlea, and hopefully improve hearing."

How did you perform the study?

"We mapped the gene expression of auditory neurons over several days of development, using single-cell transcriptomics approach, and used advanced analytical tools to reveal a hierarchical tree of neuronal diversification," says Saida Hadjab, principal researcher at the



Department of Neuroscience and co-last author of the article. "We could explore in detail, at single cell level, the dynamic of molecular developmental changes as neuron types were diverging and their identities were progressively acquired."

"We publish the whole molecular program that defines the diversification events. We then use mouse genetics to functionally evaluate this program, and eventually compare it with published datasets of deafness genes, revealing deafness genes associated with neuronal differentiation," she continues.

The next step

Several years ago, the researchers showed that neuronal diversity in the peripheral auditory system had a strong molecular basis, providing ways to manipulate each subtype to better understand their role in hearing.

With this study they demonstrate the molecular programs that set this diversity before the peripheral <u>auditory system</u> is even active. "Our next step is to explore if and how this diversity is affected in pathological conditions, for instance after noise trauma or aging, and use our new molecular knowledge and tools for resetting auditory neuron function and recovering normal hearing," says Francois Lallemend.

More information: Charles Petitpré et al, Single-cell RNA-sequencing analysis of the developing mouse inner ear identifies molecular logic of auditory neuron diversification, *Nature Communications* (2022). DOI: 10.1038/s41467-022-31580-1

Provided by Karolinska Institutet



Citation: Neurons specialized in encoding sound emerge before birth (2022, July 11) retrieved 30 June 2024 from

https://medicalxpress.com/news/2022-07-neurons-specialized-encoding-emerge-birth.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.