

New paper examines transfusion utilization and appropriateness

July 8 2022



Credit: Pixabay/CC0 Public Domain

In a new paper, researchers from University Hospitals (UH) detail how they used data via a dashboard to decrease the use of packed red-blood cell (pRBC) transfusions and platelets with an increase in appropriate



transfusions.

Their paper appears in the *Physician Leadership Journal*.

Lead author Jennifer Dawson, MBA, MSN, RN, says that they "hypothesized that use of an evidence-based <u>transfusion</u> algorithm considering patient specific criteria would raise awareness that would lead to a decrease in pRBC and platelets use, while improving appropriate use." Dawson is an operations engineer and high-reliability medicine strategist at UH.

The researchers developed a transfusion appropriateness algorithm (TAA) to analyze blood transfusions for all <u>adult patients</u> at a tertiary academic medical center. A multidisciplinary specialist panel of more than 30 participants was assembled to review current literature and existing <u>practice guidelines</u> to define clinical criteria for appropriate transfusion; with evidence suggesting that such a team approach enhances <u>patient safety</u>.

All collected data were then aggregated into a dashboard presenting the lab thresholds for each blood component alongside the appropriateness score derived from the TAA. The <u>dashboard</u> was created over the course of a year beginning in 2017, then tested and refined in 2018. It was implemented in 2019. The application housed multiple reports and allowed for reviewing the provider or patient lists for comparison or further study. A consolidated report for each patient encounter showed individual transfusion data across time, including each provider who prescribed the treatments.

The authors found a decrease in pRBC and <u>platelet</u> transfusions: 46,835 pRBC transfusions occurred in total, decreasing from 24,607 to 22,228 between groups, and 15,132 platelet transfusions, decreasing from 7,989 to 7,143 between groups. Transfusion of pRBC decreased from a mean



$2,051 (\pm 109)$ to a mean $1,852 (\pm 89)$ units per month (p =

Citation: New paper examines transfusion utilization and appropriateness (2022, July 8) retrieved 24 April 2024 from

https://medicalxpress.com/news/2022-07-paper-transfusion-appropriateness.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.