

Percentage of overdose deaths involving methadone declined between January 2019 and August 2021

July 13 2022



Credit: Pixabay/CC0 Public Domain

The percentage of methadone-involved overdose deaths relative to all drug overdose deaths declined from January 2019 to August 2021,

according to a new study. Access to methadone, a medication to treat opioid use disorder, was expanded at the start of the COVID-19 pandemic to allow more patients to take home doses, rather than visit a clinic daily. These data indicate that broader access to treatment was not associated with harms. While drug overdose deaths both with and without methadone increased in the month of March 2020, overdose deaths that did not involve methadone continued to increase in the months after the policy changes, while overdose deaths involving methadone held steady.

Published today in *JAMA Psychiatry*, this study was a [collaborative effort](#) led by researchers at the National Institute on Drug Abuse (NIDA), part of the National Institutes of Health, and the National Center for Injury Prevention and Control, part of the Centers for Disease Control and Prevention.

In 2021, provisional data from CDC estimate more than 107,000 people died of a drug [overdose](#), with 75% of those deaths involving an opioid. The overall rise in [overdose deaths](#) is largely attributable to the proliferation in the drug supply of illicit fentanyl, a highly potent synthetic opioid. A key component of the federal government response to the overdose crisis is expanding access to medications for opioid use disorder. However, only [18%](#) of people with opioid use disorder receive medication as treatment. Though the benefits of providing safe and effective medication for opioid use disorder are well-known, decades of stigma against treating [substance use disorders](#) with medication has contributed to minimal reach.

"Treatment is an essential tool to stop the addiction and overdose crises, but it is vastly underused," said NIDA Director and senior author, Nora Volkow, M.D. "This evidence adds significant weight to the argument that [effective treatment](#) for substance use disorders should be offered in an accessible and practical way that works for people who need it."

In the United States, [methadone](#) for the treatment for [opioid use disorder](#) can only be provided through federally certified opioid treatment programs, where most patients are required to visit a clinic in-person, on a daily basis, in order to get their medication. For decades, this requirement has been identified as an often-insurmountable barrier to access and retention for this treatment, particularly for people trying to balance employment, childcare, and other needs. The requirement presented unique challenges during the COVID-19 pandemic, as accessing in-person treatment became limited due to concerns about exposure to COVID-19. In order to ensure continuity of care for individuals receiving methadone treatment, on March 16, 2020, the Substance Abuse and Mental Health Services Administration (SAMHSA) [allowed states to request exceptions](#) to provide up to 28 days and 14 days of take-home methadone for stable and less stable patients, respectively.

To assess the impact and potential harms of these policy changes, investigators used data from January 2019 through August 2021 from the CDC's National Vital Statistics System, a national mortality database. Researchers calculated monthly [drug overdose](#) deaths without methadone, monthly [drug overdose deaths](#) involving methadone, and the percentage of overall overdose deaths involving methadone. They then assessed whether there was a shift in outcomes before and after the methadone take-home policy change in March 2020. They did this through interrupted time series analyses, a method of evaluation for large scale [public health interventions](#) with well-defined starting points.

Researchers found that non-methadone-involved overdose deaths increased by an average of 78 more deaths each month before March 2020, increased by 1,078 deaths during March 2020, and then continued to increase by an average of 69 more deaths each month after March 2020. Methadone-involved overdose deaths experienced a similar increase in March 2020 (increased by 94 deaths). However, the trend in

number of deaths per month before and after this initial uptick remained stable, and the percentage of overdose deaths involving methadone declined at similar rates before and after the take-home policy change, declining from 4.5% of overdose deaths in January of 2019 to 3.2% in August 2021.

Taken together, these findings indicate that the modest increase in methadone-involved overdose deaths in March 2020 was likely a reflection of the overall spike in overdose deaths driven by illicitly produced fentanyl and not an outcome of the take-home policy change. Mirroring findings from smaller studies, these national data provide evidence that the expanded opioid treatment program take-home methadone policy change established in March 2020 was not associated with increases in methadone-involved overdose deaths, despite marked increases in overall overdose deaths during the study period. Coupled with other studies that have demonstrated positive benefits related to these policies, the authors note that these findings can inform decisions about permanently expanding take-home methadone from opioid [treatment](#) programs.

"The goal of health policy should be to promote health and reduce harm, and our goal in conducting studies like this is to ensure that those policies are based on the best available scientific evidence," said lead author Christopher M. Jones, Pharm.D., Dr.P.H., acting director of the National Center for Injury Prevention and Control at the CDC. "Projects like this also underscore the important findings that can emerge when we collaborate across agencies under a common mission, as we continue to work together to address the overdose crisis."

More information: Methadone-Involved Overdose Deaths in the United States Before and After Federal Policy Changes Expanding Take-Home Methadone Doses from Opioid Treatment Programs, *JAMA Psychiatry* (2022). [DOI: 10.1001/jamapsychiatry.2022.1776](https://doi.org/10.1001/jamapsychiatry.2022.1776)

Provided by National Institutes of Health

Citation: Percentage of overdose deaths involving methadone declined between January 2019 and August 2021 (2022, July 13) retrieved 8 May 2024 from <https://medicalxpress.com/news/2022-07-percentage-overdose-deaths-involving-methadone.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.