

Periodic fasting tied to less severe COVID-19 outcomes

July 28 2022



Routine periodic fasting is associated with a lower risk for

hospitalization or mortality in patients with COVID-19, according to a study published online July 1 in *BMJ Nutrition, Prevention & Health*.

Benjamin D. Horne, Ph.D., from Intermountain Medical Center in Salt Lake City, and colleagues evaluated associations between periodic fasting and COVID-19 severity, as well as [initial infection](#) by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The analysis included 205 individuals who tested positive for SARS-CoV-2 during March 2020 to February 2021 and 1,524 with any SARS-CoV-2 test result.

The researchers found that 35.6 percent of participants engaged in periodic fasting prior to COVID-19 diagnosis. The composite outcome (COVID-19 hospitalization or mortality) occurred in 11.0 percent of periodic fasters and 28.8 percent of nonfasters. Predictors of hospitalization and mortality were age, Hispanic ethnicity, prior myocardial infarction, prior transient ischemic attack, and [renal failure](#). There was no significant association between fasting and COVID-19 diagnosis (14.3 percent in fasters and 13.0 percent in nonfasters).

"Fasting may be a [complementary therapy](#) to vaccination that could provide immune support and hyperinflammation control during and beyond the pandemic," the authors write.

More information: [Abstract/Full Text](#)

Copyright © 2022 [HealthDay](#). All rights reserved.

Citation: Periodic fasting tied to less severe COVID-19 outcomes (2022, July 28) retrieved 6 May 2024 from <https://medicalxpress.com/news/2022-07-periodic-fasting-tied-severe-covid-.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.