

Physical therapy noninferior to arthroscopic surgery for meniscal tear

July 8 2022



Exercise-based physical therapy is noninferior to arthroscopic partial

meniscectomy for patient-reported knee function at five years among patients with a degenerative meniscal tear, according to a study published online July 8 in *JAMA Network Open*.

Julia C.A. Noorduyn, from OLVG Amsterdam, and colleagues compared the five-year effectiveness of arthroscopic partial meniscectomy and [exercise](#)-based physical therapy in 321 patients aged 45 to 70 years with a degenerative meniscal tear. Participants were randomly assigned to arthroscopic partial meniscectomy or 16 sessions of exercise-based physical therapy; 87.1 percent completed the five-year follow-up (mean follow-up, 61.8 months).

The researchers found that mean improvement was 29.6 and 25.1 points on the International Knee Documentation Committee Subjective Knee Form in the [surgery](#) and physical therapy groups, respectively, from baseline to follow-up. There was a 3.5-point crude between-group difference (95 percent confidence interval, 0.7 to 6.3 [points](#); P

"Findings from this trial further support the recommendation that exercise-based [physical therapy](#) should be the preferred treatment over surgery for degenerative meniscal tear," the authors write.

More information: [Abstract/Full Text](#)
[Editorial](#)

Copyright © 2022 [HealthDay](#). All rights reserved.

Citation: Physical therapy noninferior to arthroscopic surgery for meniscal tear (2022, July 8) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-07-physical-therapy-noninferior-arthroscopic-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.