

In survey, COVID-19 vaccine recipients report changes in menstrual bleeding

July 15 2022



In a survey of more than 35,000 people, U. of I. anthropology professor Kathryn Clancy and her colleagues collected evidence that many individuals experience menstrual changes after COVID-19 vaccination. Credit: Fred Zwick, University of Illinois at Urbana-Champaign

A new analysis of reports from more than 35,000 people offers the most

comprehensive assessment so far of menstrual changes experienced by pre- and post-menopausal individuals in the first two weeks after receiving the COVID-19 vaccine. Published in the journal *Science Advances*, the study adds to the evidence that significant numbers of people experience this unexpected side effect.

"Menstruating and formerly menstruating people began sharing that they experienced unexpected bleeding after being administered a COVID-19 [vaccine](#) in early 2021," the scientists who led the study wrote. Because [vaccine trials](#) typically do not ask about [menstrual cycles](#) or bleeding, this side effect was largely ignored or dismissed.

Doctors who weighed in on the subject after hearing early reports about post-vaccination menstrual changes were often [dismissive of patients' concerns](#), said Kathryn Clancy, a professor of anthropology at the University of Illinois Urbana-Champaign who led the research with Katharine Lee, an anthropology professor at Tulane University. Some clinicians said it was unclear how a vaccine could trigger such changes.

However, other vaccines—including those for typhoid, Hepatitis B and HPV—are sometimes associated with changes in menstruation, Clancy said. These side effects are believed to be tied to an uptick in immune-related inflammatory pathways and are less likely to be driven by [hormonal changes](#).

"We suspect that for most people the changes associated with COVID-19 vaccination are short-term, and we encourage anyone who is worried to contact their doctor for further care," Lee said. "We want to reiterate that getting the vaccine is one of the best ways to prevent getting very sick with COVID, and we know that having COVID itself can lead not only to changes in periods, but also hospitalization, long COVID and death."

The researchers used a survey to query people about their experiences after vaccination. Launched in April 2021, the survey asked for demographic and other information but focused on respondents' reproductive history and experiences regarding menstrual bleeding. The team downloaded the data from the surveys on June 29, 2021. Only those who had not been diagnosed with COVID-19 were included in the analysis, as COVID-19 itself is sometimes associated with menstrual changes. The study also excluded data from people 45 to 55 years old to avoid confounding the results by including menstrual changes associated with perimenopause.

"We focused our analysis on those who regularly menstruate and those who do not currently menstruate but have in the past," Clancy said. "The latter group included postmenopausal individuals and those on hormonal therapies that suppress menstruation, for whom bleeding is especially surprising."

A [statistical analysis](#) revealed that 42.1% of menstruating survey respondents reported a heavier menstrual flow after receiving the COVID-19 vaccine. Some experienced this in the first seven days but many others saw changes 8–14 days after vaccination. Roughly the same proportion, 43.6%, reported no alteration of their menstrual flow after the vaccine, and a smaller percentage, 14.3%, saw a mix of no change or lighter flow, the researchers report.

Because the study relied on self-reported experiences logged more than 14 days after vaccination, it cannot establish causality or be seen as predictive of people in the general population, Lee said. But it can point to potential associations between a person's reproductive history, hormonal status, demographics and changes in menstruation following COVID-19 vaccination.

For example, the analysis revealed that respondents who had

experienced a pregnancy were most likely to report heavier bleeding after vaccination, with a slight increase among those who had not given birth. A majority of non-menstruating premenopausal respondents on hormonal treatment experienced breakthrough bleeding after receiving the vaccine. More than 70% of respondents using long-acting reversible contraception and 38.5% of those undergoing gender-affirming hormone treatments reported this side effect.

Participants who classified themselves as non-white, Hispanic/Latinx, or who were older, and those who experienced fever or fatigue as a side effect of vaccination were more likely than other groups to report heavier menstrual flow after vaccination. Those who had experienced endometriosis, menorrhagia, fibroids or other reproductive problems also were more likely to report a heavier menstrual flow post-vaccination, the team found.

While the uptick in menstrual flow for some people may be transitory and quickly resolve, unexpected changes in menstruation can still cause concern, Lee said.

"Unexpected breakthrough bleeding is one of the early signs of some cancers in post-menopausal people and in those who use gender-affirming hormones, so experiencing it can make people worry and require expensive and invasive cancer-screening procedures," Lee said.

"This screening is very important so we can catch cancers early," Clancy said. "Anyone with breakthrough bleeding should see their doctor. For diagnostic purposes, it would be helpful to know whether there are other causes for the bleeding."

"We'd love to see future vaccine testing protocols incorporate questions about menstruation that go beyond screening for pregnancy," Lee said. "Menstruation is a regular process that responds to all kinds of immune

and energetic stressors, and people notice changes to their bleeding patterns, yet we don't tend to talk about it publicly."

More information: Katharine M. N. Lee et al, Investigating trends in those who experience menstrual bleeding changes after SARS-CoV-2 vaccination, *Science Advances* (2022). [DOI: 10.1126/sciadv.abm7201](https://doi.org/10.1126/sciadv.abm7201). www.science.org/doi/10.1126/sciadv.abm7201

Provided by University of Illinois at Urbana-Champaign

Citation: In survey, COVID-19 vaccine recipients report changes in menstrual bleeding (2022, July 15) retrieved 18 April 2024 from <https://medicalxpress.com/news/2022-07-survey-covid-vaccine-recipients-menstrual.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.