

A type of 'step therapy' is an effective strategy for diabetic eye disease

July 14 2022



Credit: Unsplash/CC0 Public Domain

Clinical trial results from the DRCR Retina Network suggest that a specific step strategy, in which patients with diabetic macular edema start with a less expensive medicine and switch to a more expensive

medicine if vision does not improve sufficiently, gives results similar to starting off with the higher-priced drug. The main complication of diabetic macular edema, fluid build-up in the retina that causes vision loss, is commonly treated with anti-vascular endothelial growth factor (VEGF) drugs.

Results of the trial, which examined a stepped regimen of anti-VEGF drugs Avastin (bevacizumab) and Eylea (aflibercept), were published today in the *New England Journal of Medicine*.

"Our study showed that switching treatments when needed is a reasonable strategy," said Chirag Jhaveri, M.D., Austin Research Center for Retina, Texas, the lead study author. "Insurance companies often require clinicians to start with the less expensive treatment, so we really wanted to see how a specific treatment strategy using this approach would affect patient care."

Diabetic macular edema is caused by diabetes-related alterations to retinal blood vessels. Symptoms include blurred vision. If untreated, [vision loss](#) can become permanent and progress to blindness. Retinal injections of anti-VEGF drugs can restore vision. The DRCR Retina Network previously showed that Avastin and Eylea improve visual acuity in people with [diabetic macular edema](#). However, while Eylea is approved by the U.S. Food and Drug Administration to treat diabetic macular edema and results in better visual outcomes on average, off-label Avastin is much less expensive and is sometimes required by insurers as a first-line treatment.

The study enrolled 270 participants with diabetic macular edema, some of whom received treatments in both eyes. At enrollment, all had best-corrected visual acuity between 20/50 and 20/320. Half the study eyes were assigned to Eylea from the start, and half were assigned to start with Avastin. For participants who needed treatment in both eyes, each

eye started treatment with a different drug. Participants received either Avastin or Eylea injections every four weeks for 24 weeks. If eyes assigned Avastin failed to reach the pre-set improvement benchmarks starting at 12 weeks, the eye was switched to Eylea.

After 24 weeks, physicians could taper down the frequency of injections as appropriate to maintain visual acuity. The study collected information about participants' retinal structure and visual acuity for two years.

After two years, eyes in both groups had similar [visual acuity](#) outcomes, improving on average approximately three lines on an eye chart, compared to the trial's start. In the Avastin group, 70% of eyes switched to Eylea during the study.

"While most participants on Avastin eventually switched to Eylea, they still had improvement during those initial weeks, even if they didn't hit our pre-set benchmarks," said Adam Glassman of the Jaeb Center for Health Research and director of the DRCR Retina Network coordinating center. "There are large cost disparities between these drugs, so differences in treatment strategies may have substantial cost implications."

"We've demonstrated here one method to managing a step treatment, where the outcomes are similar to the best existing treatment protocol with Eylea," said Jennifer Sun, M.D., M.P.H., of Joslin Diabetes Center and Harvard Medical School, Boston, and chair of diabetes initiatives for the DRCR Retina Network. "Any time we can add to a clinician's toolbox, whether it's a new medication or a new approach to using existing medications, as in this study, it's a benefit for patients."

More information: Jhaveri CD et al, Aflibercept Monotherapy or Bevacizumab First for Diabetic Macular Edema, *New England Journal of Medicine* (2022). [DOI: 10.1056/NEJMoa2204225](https://doi.org/10.1056/NEJMoa2204225)

Provided by National Eye Institute

Citation: A type of 'step therapy' is an effective strategy for diabetic eye disease (2022, July 14) retrieved 27 June 2024 from <https://medicalxpress.com/news/2022-07-therapy-effective-strategy-diabetic-eye.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.