

Women urged to eat potassium-rich foods to improve their heart health

July 22 2022



To analyze whether the association between daily potassium consumption and systolic blood pressure (SBP) as well as cardiovascular (CV) events differs between men and women.

To analyze whether these associations are present with increasing amounts of sodium intake.

Key Finding


Interaction by sex was observed for the association between potassium intake, SBP and CV events.

In women, but not in men, the inverse slope between potassium intake and SBP was steeper in those with a higher sodium intake.

Take Home Message

This study demonstrates that the association between potassium intake and both SBP and CV events is sex-specific.

The data suggest that particularly high sodium-consuming women may benefit from a high potassium intake with regards to SBP.





An analysis was performed in 11 267 men and 13 696 women from the EPIC-Norfolk cohort. Twenty-four hour excretion of sodium and potassium, reflecting intake, was estimated from sodium and potassium concentration in spot urine samples using the Kawasaki formula. Linear and Cox regression analyses showed that the association between potassium intake and both SBP and CVD risk is sex specific. Moreover, among women, the association between potassium intake and SBP was modified by sodium intake. EPIC-Norfolk, Norfolk cohort of the European Prospective Investigation into Cancer; CVD, cardiovascular disease; SBP, systolic blood pressure. Credit: *European Heart Journal* (2022). DOI: 10.1093/eurheartj/ehac313

Women who eat bananas, avocados and salmon could reduce the negative effects of salt in the diet, according to a study published today in the *European Heart Journal*, a journal of the European Society of Cardiology (ESC). The study found that potassium-rich diets were associated with lower blood pressure, particularly in women with high salt intake.

"It is well known that high salt consumption is associated with elevated blood pressure and a raised risk of heart attacks and strokes," said study author Professor Liffert Vogt of Amsterdam University Medical Centers, the Netherlands. "Health advice has focused on limiting salt intake but this is difficult to achieve when our diets include processed foods. Potassium helps the body excrete more sodium in the urine. In our study, dietary potassium was linked with the greatest health gains in women."

The study included 24,963 participants (11,267 men and 13,696 women) of the EPIC-Norfolk study, which recruited 40- to 79-year-olds from general practices in Norfolk, U.K., between 1993 and 1997. The average age was 59 years for men and 58 years for women. Participants completed a questionnaire on lifestyle habits, blood pressure was



measured, and a urine sample was collected. Urinary sodium and potassium were used to estimate <u>dietary intake</u>. Participants were divided into tertiles according to sodium intake (low/medium/high) and <u>potassium intake</u> (low/medium/high).

The researchers analyzed the association between potassium intake and blood pressure after adjusting for age, sex and sodium intake. Potassium consumption (in grams per day) was associated with blood pressure in women—as intake went up, blood pressure went down. When the association was analyzed according to sodium intake (low/medium/high), the relationship between potassium and blood pressure was only observed in women with high sodium intake, where every 1 gram increase in daily potassium was associated with a 2.4 mmHg lower systolic blood pressure. In men, there was no association between potassium and blood pressure.

During a median follow-up of 19.5 years, 13,596 (55%) participants were hospitalized or died due to <u>cardiovascular disease</u>. The researchers analyzed the association between potassium intake and <u>cardiovascular events</u> after adjusting for age, sex, body mass index, sodium intake, use of lipid lowering drugs, smoking, alcohol intake, diabetes and prior heart attack or stroke. In the overall cohort, people in the highest tertile of potassium intake had a 13% lower risk of cardiovascular events compared to those in the lowest tertile. When men and women were analyzed separately, the corresponding risk reductions were 7% and 11%, respectively. The amount of salt in the diet did not influence the relationship between potassium and cardiovascular events in men or women.

Professor Vogt said: "The results suggest that potassium helps preserve heart health, but that <u>women</u> benefit more than men. The relationship between potassium and cardiovascular events was the same regardless of <u>salt intake</u>, suggesting that potassium has other ways of protecting the



heart on top of increasing sodium excretion."

The World Health Organization recommends that adults consume at least 3.5 grams of potassium and less than 2 grams of sodium (5 grams of salt) per day. High potassium foods include vegetables, fruit, nuts, beans, dairy products and fish. For example, a 115 gram banana has 375 mg of potassium, 154 grams of cooked salmon has 780 mg, a 136 gram potato has 500 mg, and 1 cup of milk has 375 mg.

Professor Vogt concluded: "Our findings indicate that a heart healthy diet goes beyond limiting salt to boosting potassium content. Food companies can help by swapping standard sodium-based salt for a potassium salt alternative in processed foods. On top of that, we should all prioritize fresh, unprocessed foods since they are both rich in potassium and low in salt."

More information: Rosa D Wouda et al, Sex-specific associations between potassium intake, blood pressure, and cardiovascular outcomes: the EPIC-Norfolk study, *European Heart Journal* (2022). DOI: 10.1093/eurheartj/ehac313

Provided by European Society of Cardiology

Citation: Women urged to eat potassium-rich foods to improve their heart health (2022, July 22) retrieved 12 September 2024 from

https://medicalxpress.com/news/2022-07-women-urged-potassium-rich-foods-heart.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.