

# New study shows promising alternative to opioids for pain after tooth extractions

August 18 2022, by Karen Black

---



Credit: Pixabay/CC0 Public Domain

As the opioid epidemic rages on during the COVID-19 pandemic, a study by the University of Rochester Medical Center's Eastman Institute for Oral Health (EIOH) shows promise for a non-opioid alternative for

acute dental pain.

Prescription opioid-involved death rates increased more than 16% from 2019 to 2020, and among all the opioid overdose deaths, [prescription opioids](#) represent nearly 18 percent, according to the Centers for Disease Control and Prevention.

Published today in *JAMA Network Open*, the study compares two large groups of patients representing an equal number of men and women ages 18-93 who had dental [extractions](#) at the Institute's Howitt Urgent Dental Care clinic in Rochester, New York.

"We hypothesized that using a combination of the non-opioid pain medications and adding gabapentin to the mix for pain would be an effective strategy to minimize or eliminate opioids for dental pain," said Yanfang Ren, DDS, Ph.D., MPH, professor and clinical chief, Howitt Urgent Dental Care.

For the 3,300 patients in the first group in 2012, those with mild pain were treated with ibuprofen or [acetaminophen](#). For moderate to severe pain, patients were prescribed higher doses of ibuprofen or opioid combinations including hydrocodone, oxycodone or codeine.

But the second group of nearly 3,800 people who had extractions from March 2021 through February 2022, received no opioids whatsoever.

Similarly, for mild pain, the second group was prescribed acetaminophen or ibuprofen. For moderate to severe pain, higher doses of ibuprofen or an ibuprofen and acetaminophen combination were prescribed.

For the segment of patients who were unable to take ibuprofen or acetaminophen due to health issues or potential interactions with existing medications, they were often prescribed an opioid combination in year

2012, but a [gabapentin](#) combination was used instead in 2022 to use as a non-opioid alternative. The results were promising when the effectiveness was evaluated using a real-world measure of proportion of patients returning for additional pain treatment after receiving the prescribed analgesics.

"Although the American Dental Association recommends nonsteroidal anti-inflammatory drugs for managing pain," Dr. Ren said, "dentists frequently prescribe opioids for dental pain and contribute substantially to new and persistent opioid use. This may partly be explained by the lack of alternatives to opioids, especially when the [ibuprofen](#) or acetaminophen are ineffective or can't be tolerated."

"This study represents continued efforts by our team and other dentists to minimize the use of [opioids](#) for [dental pain](#)," said Eli Eliav, DMD, Ph.D., the director of EIOH and a pain expert. "Additional studies, preferably randomized controlled clinical trials, are needed to confirm the safety and effectiveness of this approach. It is our duty to continuously seek safe and effective treatment for our patients in pain."

Other authors on the study include Qirong Huang, DDS, MS, Linda Rasubala, DDS, Ph.D., Richard H. Gracely, Ph.D., Junad Khan, BDS, MSD, MPH, Ph.D., and Eli Eliav, DMD, Ph.D.

**More information:** Qirong Huang et al, Comparison of Analgesic Prescriptions for Dental Pain and Patient Pain Outcomes Before vs After an Opioid Reduction Initiative, *JAMA Network Open* (2022). [DOI: 10.1001/jamanetworkopen.2022.27219](https://doi.org/10.1001/jamanetworkopen.2022.27219)

Provided by University of Rochester Medical Center

Citation: New study shows promising alternative to opioids for pain after tooth extractions (2022, August 18) retrieved 18 April 2024 from <https://medicalxpress.com/news/2022-08-alternative-opioids-pain-tooth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.