

## **Study shows PPE was highly effective against COVID-19 in emergency department workers**

August 9 2022, by Bob Shepard



Credit: Unsplash/CC0 Public Domain

Workers in America's emergency departments were more likely to contract COVID-19 in the early days of the pandemic in community settings or at home rather than when providing patient care in a hospital



setting, according to new research conducted at emergency departments across the nation.

The study, part of the COVID-19 Evaluation of Risks in Emergency Departments, known as the Project COVERED study, found that <u>personal protective equipment</u>, including masks, gloves and protective gowns, helped keep emergency health care workers shielded from infection while caring for patients. The study was published July 22 in the journal *PLOS ONE*.

"From the start of the pandemic, emergency department personnel served as the initial <u>hospital</u> caregivers for the majority of critically ill patients with known or suspected COVID-19 infection," said Walter Schrading, M.D., professor in the Department of Emergency Medicine in the Heersink School of Medicine at the University of Alabama at Birmingham. Schrading is a co-author on the study.

Between May and December 2020, prior to the COVID-19 vaccine availability, emergency department staff at 20 medical centers across the country were followed with surveys, COVID-19 tests and serology testing. The study included 1,673 emergency department physicians, advanced practice providers, nurses and non-clinical staff who were observed over 20 weeks and 30,000 person-weeks in total. During this surveillance, more than 4,400 high-risk aerosolizing intubations were performed.

"At the beginning of the pandemic, the risk of providing emergency patient care was not understood. Emergency care staff were committed to their role in responding to the crisis, but they were understandably anxious," said Nicholas Mohr, M.D., lead author and co-principal investigator of the study, at the University of Iowa Hospitals and Clinics. "Even though emergency care workers felt vulnerable performing procedures and providing clinical care for COVID-19 patients during



this time, our findings support that it was really when they were outside the hospital and at home, with their families and friends, and in their community that people were at risk of infection."

The study found that only 4.5% of <u>emergency personnel</u> became infected with COVID-19, and the most prominent factors associated with those infections were community exposures, hospital case counts and mask use in public.

"This study does provide reassurance to <u>health care professionals</u> that, with the proper precautions, they can be relatively safe while treating patients in the emergency medicine setting," Schrading said. "PPE is effective at preventing COVID-19 transmission during <u>clinical care</u>, so when case counts are high in the community, health care personnel should take precautions outside the hospital just as they do inside the hospital when they are working."

**More information:** Nicholas M. Mohr et al, Emergency department personnel patient care-related COVID-19 risk, *PLOS ONE* (2022). DOI: 10.1371/journal.pone.0271597

Provided by University of Alabama at Birmingham

Citation: Study shows PPE was highly effective against COVID-19 in emergency department workers (2022, August 9) retrieved 14 May 2024 from https://medicalxpress.com/news/2022-08-ppe-highly-effective-covid-emergency.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.