

# Proximity to fracking sites associated with risk of childhood cancer

August 17 2022

---



Credit: Pixabay/CC0 Public Domain

Pennsylvania children living near unconventional oil and gas (UOG) developments at birth were two to three times more likely to be diagnosed with leukemia between the ages of 2 and 7 than those who did

not live near this oil and gas activity, after accounting for other factors that could influence cancer risk, a novel study from the Yale School of Public Health finds.

The registry-based study, published Aug. 17 in the journal *Environmental Health Perspectives*, included nearly 2,500 Pennsylvania children, 405 of whom were diagnosed with [acute lymphoblastic leukemia](#), the most common type of [cancer](#) in children.

Acute lymphoblastic leukemia, also referred to as ALL, is a type of leukemia that arises from mutations to lymphoid immune cells. Although long-term survival rates are high, children who survive this disease may be at higher risk of other [health](#) problems, developmental challenges, and psychological issues. Unconventional oil and gas development, more commonly referred to as fracking (short for [hydraulic fracturing](#)), is a method for extracting gas and oil from shale rock. The process involves injecting water, sand, and chemicals into bedrock at high pressure, which allows gas and oil to flow into a well and then be collected for market.

For communities living nearby, UOG development can pose a number of potential threats. Chemical threats include, for example, air pollution from vehicle emissions and well and road construction, and [water pollution](#) from hydraulic fracturing or spills of wastewater. Hundreds of chemicals have been reportedly used in UOG injection water or detected in wastewater, some of which are known or suspected to be cancer causing. The paucity of data on the association between UOG and childhood cancer outcomes has fueled [public concerns](#) about possible cancer clusters in heavily drilled regions and calls for more research and government action.

"Unconventional oil and gas development can both use and release chemicals that have been linked to cancer, so the potential for children

living near UOG to be exposed to these chemical carcinogens is a major public health concern," said the study's senior author, Nicole Deziel, associate professor of epidemiology at the Yale School of Public Health.

"Studies of UOG exposure and cancer are extremely few in number. We set out to conduct a high-quality study to further investigate this potential relationship," added Cassandra Clark, the study's first author and a postdoctoral associate at the Yale Cancer Center. "Our results indicate that exposure to UOG may be an important risk factor for ALL, particularly for children exposed in utero."

The study also found that [drinking water](#) could be an important pathway of exposure to oil and gas-related chemicals. The authors applied a new exposure metric in this study that they call "IDups" (which stands for "inverse distance to the nearest upgradient unconventional oil and gas well"). This means that the researchers identified UOG wells that fell within a child's watershed area—the zone from which a drinking water well serving their home would likely draw water—and calculated the distance from the home to the nearest of those UOG wells. UOG wells falling within the watershed area are expected to be more likely to impact the home's drinking water supply, they said.

"Previous health studies have found links between proximity to oil and gas drilling and various children's health outcomes," said Deziel. "This study is among the few to focus on drinking water specifically and the first to apply a novel metric designed to capture potential exposure through this pathway."

This work adds to a growing body of literature on UOG exposure and children's health used to inform policy, such as setback distances (the required minimum distance between a private residence or other sensitive location and a UOG well). Current setback distances are the subject of much debate in the United States, with some calling for

setback distances to be lengthened to more than 305 meters (1,000 feet) and as far as 1,000 meters (3,281 feet). The allowable setback in Pennsylvania, where the study was conducted, is 500 feet, or 152 meters.

"Our findings of increased risk of ALL at distances of two kilometers or more from UOG operations, in conjunction with evidence from numerous other studies, suggest that existing setback distances, which may be as little as 150 feet, are insufficiently protective of children's health," Clark said. "We hope that studies like ours are taken into account in the ongoing policy discussion around UOG setback distances."

**More information:** Cassandra J. Clark et al, Unconventional Oil and Gas Development Exposure and Risk of Childhood Acute Lymphoblastic Leukemia: A Case–Control Study in Pennsylvania, 2009–2017, *Environmental Health Perspectives* (2022). [DOI: 10.1289/EHP11092](https://doi.org/10.1289/EHP11092)

Provided by Yale University

Citation: Proximity to fracking sites associated with risk of childhood cancer (2022, August 17) retrieved 28 April 2024 from

<https://medicalxpress.com/news/2022-08-proximity-fracking-sites-childhood-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.