


# Reprogramming human skin cells into aged neurons to study neurodegenerative disorders

August 17 2022



HD-iNs exhibit neurite specific autophagy alteration. (A) Reduced BECN1 expression in HD-iNs compared to Ctrl-iNs using western blot (n = 10 replicates for control and n = 9 replicates for HD-iNs). (B) LC3B-II levels are significantly increased in the HD-iNs, while the LC3B-II/I ratio decreased compared to the healthy Ctrl-iNs (n = 6 replicates). (C–H) Representative images and statistical analysis shows a significant increase both in number and size of LC3B, p62 and LAMP1 dots in the MAP2<sup>+</sup> neurites of HD-iNs compared to controls (n = 6 lines). (I) Representative images of human post-mortem striatal tissue from a

healthy control and three different Huntington's disease patients at different disease stages showing p62 accumulation specifically in the neurites as visualized by a neurofilament specific antibody. \*\*\*P

Citation: Reprogramming human skin cells into aged neurons to study neurodegenerative disorders (2022, August 17) retrieved 6 May 2024 from <https://medicalxpress.com/news/2022-08-reprogramming-human-skin-cells-aged.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.