

Modeling the use of beta radiation in cancer treatment

September 21 2022




Illustration of beta decay proceeding against the backdrop of a Monte Carlo simulation. Credit: Robert Lea

Treating superficial skin tumors especially when they are located above cartilage or bone with beta radiation can help protect sensitive structures during the delivery of treatment.

The use of short-range beta radiation in [cancer treatment](#) is not without its disadvantages, however, especially when it comes to the measurement of radiation exposure — dosimetry. When experimental dosimetry is not feasible, researchers use simulations and calculations to study the interaction of the ionizing radiation with matter and estimate the radiation dose delivered to a target organ.

A new paper published in *The European Physical Journal Plus* and authored by Eduardo De Paiva, from the Division of Medical Physics at the Institute of Radiation Protection and Dosimetry, Rio de Janeiro, Brazil, and his colleagues, pits the gold standard of [simulation](#) techniques — Monte Carlo (MC) simulation — against an alternative analytic method, the Loevinger formula.

"We measured the dose of a treatment applicator using mathematical techniques — a simple technique, no experiment needed and no practical challenges," De Paiva says. "The comparison of MC simulation and Loevinger formula on the setup of our research was the novelty of our study."

Nonexperimental dosimetry techniques like MC simulation are advantageous for their ability to handle different geometries and materials, but MC simulations require heavy computation and this can impede their implementation.

Analytic methods are another set of techniques for dosimetry of beta radiation that can produce results faster than MC methods. Thus far, these methods have been less favored because they are associated with lower accuracy.

The team used MC simulation and analytical calculation — the Loevinger formula — for dosimetry of [radiation dose](#) from a multi-well skin brachytherapy applicator with two beta sources. The results of the

two approaches were compared to see how accurate the [analytical method](#) is.

"The Loevinger formula, which is a quick method for dosimetry showed a good agreement with gold standard Monte Carlo methods," Paiva concluded. "Thus, the Loevinger formula can be used, as the basis of a dosimetry software, for straightforward dosimetry of beta sources in simple geometries."

More information: Eduardo De Paiva et al, Calculations of beta radiation doses from multiwell Phosphorus-32 and Yttrium-90 applicators designed to be used in the treatment of superficial skin tumors: comparison of Monte Carlo and analytical methods, *The European Physical Journal Plus* (2022). [DOI: 10.1140/epjp/s13360-022-03116-5](#)

Provided by Springer

Citation: Modeling the use of beta radiation in cancer treatment (2022, September 21) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-09-beta-cancer-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--