

Bronchodilators don't improve smokingrelated respiratory symptoms in people without COPD

September 4 2022



Credit: Pixabay/CC0 Public Domain

Researchers supported by the National Institutes of Health have found that dual bronchodilators—long-lasting inhalers that relax the airways



and make it easier to breathe—do little to help people who do not have chronic obstructive pulmonary disease (COPD), but who do have respiratory symptoms and a history of smoking.

COPD, a <u>lung disease</u> that obstructs the airways and leads to coughing, wheezing, and shortness of breath, affects about 15 million Americans. However, millions of others who smoke or used to smoke and have some symptoms of COPD have also been prescribed bronchodilators.

"We've assumed these medications worked in <u>patients</u> who don't meet lung function criteria for COPD, but we never checked," said MeiLan K. Han, M.D., a principal investigator and first author of the study. "We now know these existing medications don't work for these patients."

The findings of the study, which was funded by the National Heart, Lung, and Blood Institute (NHLBI), were published in the *New England Journal of Medicine* and simultaneously presented at the European Respiratory Society International Congress.

According to scientists, the implications are significant. First, they show the importance of diagnosing lung conditions through spirometry, a lung function test Han noted is underutilized in clinical practice. Second, they show the need for new, <u>effective therapies</u> for patients without COPD.

Inhalers have long been the primary go-to treatment for these patients, she explained, because doctors either assume a patient has COPD, or if they don't, that their smoking-related symptoms could be helped by the inhalers. But while tobacco smoking causes a large spectrum of lung damage, the study showed bronchodilator therapy only helps patients with enough lung damage that would result in abnormal spirometry readings.

In the 12-week, randomized, double-blinded study, which was part of the



Redefining Therapy in Early COPD for the Pulmonary Trials Cooperative (RETHINC), researchers enrolled 535 adults with symptoms of COPD, ages 40-80, at one of 20 U.S. medical centers. Twice each day, study participants used an inhaler that contained either medication or a placebo.

By the end of the trial, some adults in the medication (intervention) and placebo (control) groups saw slight respiratory improvements—this could mean they coughed less, produced less phlegm, or felt less winded—which was assessed through the <u>St. George's Respiratory</u> <u>Questionnaire</u>. However, the researchers found no significant differences between those receiving medication or placebo. They reported 56% (128 of 227) of participants who received the medication saw respiratory symptom improvements, compared to 59% (144 of 244) of those who took the placebo.

According to Han, these data underscore why we can't continue to do what we have been doing, which is not doing spirometry and just treating patients with the same COPD medications and expect that we're going to see improvement.

Antonello Punturieri, M.D., Ph.D., program director of NHLBI's Chronic Obstructive Pulmonary Disease/Environment Program, said spirometry testing should be used for any patient who shows signs of COPD, airflow obstruction, or who has a history of cigarette smoking. Though spirometry readings are used during about <u>one-third</u> of medical visits related to COPD, roughly half of patients who would meet criteria for COPD go undiagnosed.

Helping patients quit smoking is a primary way to prevent COPD or COPD-like symptoms, the study noted. More than <u>30 million</u> adults smoke, according to the Centers for Disease Control and Prevention, and many who are not diagnosed with COPD share symptoms. About <u>one in</u>



four current or former smokers without COPD have reported having shortness of breath. In addition to encouraging smoking cessation, doctors can help patients who do not meet lung-function criteria of COPD by working with them to address any other underlying issues, such as overweight and obesity, <u>heart failure</u>, or other <u>lung</u> issues. "In the meantime, research should be focused on finding new treatments for them," Han explained. "The next question is, can we develop more targeted therapies for these patients who are on the milder end of the spectrum?"

"Because cough and mucus production show up prominently among these patients, we believe therapies that target <u>mucus production</u> in the airways may be effective," said Prescott G. Woodruff, M.D., a principal investigator and senior author of the study.

More information: Bronchodilators in symptomatic tobacco-exposed persons with preserved spirometry for the RETHINC study group, *New England Journal of Medicine* (2022). DOI: 10.1056/NEJMoa2204752

Provided by NIH/National Heart, Lung and Blood Institute

Citation: Bronchodilators don't improve smoking-related respiratory symptoms in people without COPD (2022, September 4) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2022-09-bronchodilators-dont-smoking-related-respiratory-symptoms.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.