

Study identifies emerging illnesses from COVID-19 in children

September 29 2022



Credit: Pixabay/CC0 Public Domain

A National Jewish Health led study finds that children who develop shortness of breath after COVID-19 infection have evidence of peripheral airway obstruction. These children can experience persistent fatigue, shortness of breath, chest pain or cough after recovering from

their acute infection symptoms.

These symptoms, also known as "long COVID," can last for months and significantly affect the quality of life by impacting cognitive function and stamina in school, while also limiting participation in sports and everyday activities. The study was published today in the *Journal of Allergy and Clinical Immunology in Practice*.

"Many [young patients](#) who continue to experience respiratory symptoms months after having been diagnosed with COVID are told it's all in their head, because spirometry and other diagnostics fail to show any abnormalities," said Nathan Rabinovitch, MD, MPH, lead author of the study and director of the pediatric care unit at National Jewish Health as well as Professor in the Division of Pediatric Allergy and Clinical Immunology.

Peripheral [airway obstruction](#) means that small airways less than 2 mm in diameter are narrower than expected. This could be due to inflammation in the airways or constriction of the airway muscles in that area. Using sophisticated testing, physicians at National Jewish Health identified subtle changes in the small airways that likely caused respiratory symptoms.

One patient referenced in the study had no previous history of asthma and exhibited normal spirometry test results. However, he demonstrated abnormal lung hyperinflation and peripheral airway obstruction that reversed with albuterol. Over the next few months, his symptoms and airway obstruction improved with daily use of an inhaled corticosteroid, bronchodilator medication and a focused physical therapy regimen.

"Through this research, we are identifying the specific diseases that long COVID may be causing, and, therefore, we can target a specific treatment for these individual issues," said Dr. Rabinovitch.

More information: Nathan Rabinovitch et al, Reversible peripheral airway obstruction and lung hyperinflation in children presenting with dyspnea and exercise intolerance after COVID-19 infection, *The Journal of Allergy and Clinical Immunology: In Practice* (2022). [DOI: 10.1016/j.jaip.2022.07.038](https://doi.org/10.1016/j.jaip.2022.07.038)

Provided by National Jewish Health

Citation: Study identifies emerging illnesses from COVID-19 in children (2022, September 29) retrieved 23 April 2024 from <https://medicalxpress.com/news/2022-09-emerging-illnesses-covid-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.