

Measuring changes to the male brain after fathering a child

September 13 2022, by Bob Yirka



Credit: Pixabay/CC0 Public Domain

A team of researchers with members from Instituto de Investigación Sanitaria Gregorio Marañón, the University of Southern California and the Autonomous University of Barcelona has found that some parts of

the male brain change in size after a man fathers a child. In their paper published in the journal *Cerebral Cortex*, the group describes how they studied MRI scans of new fathers to learn more about how their brains might change as they adapt to fatherhood.

Many studies have looked at changes to the female brain after a woman gives birth and begins to raise a child, but little research has been done on changes to the male brain. To rectify that situation, the researchers carried out a study that involved conducting MRI scans of two groups of men taken before the birth of their child and after. The first group was in Spain—the researchers scanned each volunteer's brain before their partner became pregnant and then again shortly after the couple had a child together. The other group was in the U.S., where researchers scanned the male volunteers' brains during their partner's late pregnancy and then again approximately eight months after the baby was born. For their [control group](#), the researchers also scanned the brains of 18 male volunteers who did not sire children.

The researchers found little to no change in the [limbic system](#), which is where women tend to have the biggest changes after giving birth. But there were changes in the men's cortical gray matter, which prior research has shown plays a role in [social skills](#). The loss of volume in this region was found in what is known as the default mode network, which prior research has suggested plays a role in parental acceptance and warmth. A loss of volume, the researchers note, suggests the brain region is becoming more dense, which makes it more efficient.

The team also found that men tend to experience reductions in volume of the part of their brains that processes [visual information](#). The researchers suggest this indicates that the male [brain](#) changes to assist in recognizing their new child and responding to it.

More information: Magdalena Martínez-García et al, First-time

fathers show longitudinal gray matter cortical volume reductions: evidence from two international samples, *Cerebral Cortex* (2022). DOI: [10.1093/cercor/bhac333](https://doi.org/10.1093/cercor/bhac333)

© 2022 Science X Network

Citation: Measuring changes to the male brain after fathering a child (2022, September 13) retrieved 9 May 2024 from <https://medicalxpress.com/news/2022-09-male-brain-fathering-child.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.