


This molecule could be behind liver fibrosis

September 26 2022, by Kelly Malcom



Chronic liver disease is associated with NIK upregulation in cholangiocytes. **A, B** Human liver sections were stained with antibodies to NIK and K19. **A** Representative images. Scale bar: 200 μ m. **B** NIK⁺, K19⁺, and NIK⁺K19⁺ cells were counted and normalized to total cells. Control: *n* = 3 subjects, PBC: *n* = 3 subjects, PSC: *n* = 3 subjects, HBV: *n* = 5 subjects, HCV: *n* = 3 subjects, Alcohol: *n* = 3 subjects. **C, D** C57BL/6J male mice were fed a chow or DDC diet for 4 weeks. **C** Liver sections were stained with antibodies to NIK and K19. NIK⁺, K19⁺, and NIK⁺K19⁺ cells were counted and normalized to total cells. Chow: *n* = 3 mice, DDC: *n* = 3 mice. Scale bar: 200 μ m. **D** Liver NIK expression was measured by qPCR (normalized to 18 S levels). Chow: *n* = 4 mice, DDC: *n*

= 6 mice. a.u. arbitrary units. **E**, **F** C57BL/6J males were treated with BDL or sham surgery for 7 days. **E** Liver sections were stained with antibodies to NIK and K19. NIK⁺, K19⁺, and NIK⁺K19⁺ cells were counted and normalized to total cells ($n = 3$ mice per group). Scale bar: 200 μm . **F** Liver NIK expression was measured by qPCR (normalized to 36B4 levels, $n = 4$ mice per group). Data are presented as mean \pm SEM. * p

Citation: This molecule could be behind liver fibrosis (2022, September 26) retrieved 29 April 2024 from <https://medicalxpress.com/news/2022-09-molecule-liver-fibrosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.