


MRI findings in opioid-exposed fetuses show smaller brain size

September 28 2022



Sagittal (A, B) and coronal (C-F) T2-weighted SSFSE images show 2D measurements of brain fronto-occipital diameter (A), bone fronto-occipital diameter (B), brain biparietal diameter (C), bone biparietal diameter (D), right cerebral biparietal diameter (E), left cerebral biparietal diameter (F). Credit: ARRS/*American Journal of Roentgenology*

According to an open-access article in the *American Journal of Roentgenology (AJR)*, third-trimester fetuses with in utero opioid exposure exhibited multiple smaller 2D biometric measurements of the brain, as well as altered fetal physiology, on investigational MRI.

Noting the scarcity of imaging literature evaluating prenatal opioid exposure on [brain](#) development, "Our findings demonstrate the impact of prenatal opioid exposure on [fetal brain development](#), which may in turn affect postnatal clinical outcomes," wrote first author Usha D. Nagaraj, MD, from Cincinnati Children's Hospital Medical Center in Ohio.

Nagaraj and colleagues' prospective multicenter case-control study included 65 women (mean age, 29 years) in their third trimester of pregnancy who underwent fetal MRI from July 1, 2020 through December 31, 2021 at three U.S. academic medical centers: Cincinnati Children's Hospital Medical Center, Arkansas Children's Hospital, and the University of North Carolina at Chapel Hill. A total of 28 fetuses (mean gestational age, 32.3 weeks) were classified as opioid-exposed, while 37 fetuses (mean gestational age, 31.9 weeks) were unexposed in utero. Fourteen 2D biometric measurements of the fetal brain were manually assessed and used to derive four indices.

When adjusting for [gestational age](#), fetal sex, and nicotine exposure, seven of the 14 2D biometric measurements—cerebral fronto-occipital diameter, bone biparietal diameter, brain biparietal diameter, corpus callosum length, vermis height, anterior-posterior pons measurement, and transverse cerebellar diameter—were significantly smaller in opioid-exposed fetuses than in unexposed fetuses, as measured on fetal MRI.

"In addition," the authors of this *AJR* article concluded, "fetuses with prenatal [opioid](#) exposure had increased frequencies of breech presentation and of increased amniotic fluid volume."

More information: Usha D. Nagaraj et al, MRI Findings in Third-Trimester Opioid-Exposed Fetuses With Focus on Brain Measurements: A Prospective Multicenter Case-Control Study, *American Journal of Roentgenology* (2022). [DOI: 10.2214/AJR.22.28357](https://doi.org/10.2214/AJR.22.28357)

Provided by American Roentgen Ray Society

Citation: MRI findings in opioid-exposed fetuses show smaller brain size (2022, September 28) retrieved 11 May 2024 from <https://medicalxpress.com/news/2022-09-mri-opioid-exposed-fetuses-smaller-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.