

# Restrictions on paracetamol may be coming. What does that mean for consumers?

September 29 2022, by Natasa Gisev and Ria Hopkins

---



Credit: AI-generated image ([disclaimer](#))

Paracetamol is Australia's most widely used pain medicine, with [65 million packs sold](#) across the country in 2021. It is available everywhere from toilet vending machines, convenience stores, supermarkets, and pharmacies.

The Therapeutic Goods Administration (TGA) is now undertaking a [public consultation](#) into access to [paracetamol](#) in the community due to concerns about poisonings, especially among young people.

Adjunct Professor John Skerritt and Professor Robyn Langham gave a presentation on the key findings of the independent expert report on the risks of intentional self-poisoning with paracetamol and possible next steps.

Watch here: <https://t.co/vJbie1ozjn> [pic.twitter.com/orflsq4KyU](https://pic.twitter.com/orflsq4KyU)

— TGA Australia (@TGAgovau) [September 26, 2022](#)

## What is paracetamol and what is it used for?

Paracetamol (commonly marketed as Panadol, Panamax or Dymadon) is a [medicine used](#) to [treat pain](#) and reduce fever.

Outside of pharmacies, paracetamol is [available in packs of up to 20 tablets](#) (or capsules), while pharmacies stock [packs](#) of up to 100 tablets and a range of formulations, including liquids and suppositories. Paracetamol's wide availability and low cost make it a convenient option for people to self-treat pain without a doctor's appointment or prescription.

At [therapeutic doses](#), paracetamol is [considered safe for most people](#), with few side effects when used as directed.

However, it can be dangerous at high doses, resulting in [acute liver toxicity](#), which in severe cases may lead to death. Although there are [treatments to reverse paracetamol overdose](#), they need to be given within 2-8 hours to be most effective.

## Why is the TGA reviewing paracetamol access?

The TGA is concerned about harms from paracetamol poisoning, particularly intentional overdoses among young people, which have been seen in [Australia](#) as well as [internationally](#). In response, the TGA requested an [independent expert report](#) to support its review of paracetamol access.

The report found that between 2007 and 2020, there were 40–50 deaths each year in Australia from paracetamol poisoning. From 2009 to 2017, hospital admissions due to paracetamol poisoning increased from 8,617 to 11,697. They decreased to 8,723 in 2019–20.

Around 80% of these admissions were due to intentional self-poisoning, with young people aged between 10–24 years accounting for 40–50% of these incidents. Hospitalizations among [young people](#) due to intentional overdoses from any medicine also increased over this period. Calls to Australian Poisons Information Centers about self-poisonings involving paracetamol also increased over the last decade.

It is important to note that the number of harmful events is low compared to the amount of paracetamol sold in Australia. For every million packs of paracetamol sold, there were 100 hospital admissions for intentional self-poisoning, three [hospital admissions](#) for liver injury, and less than one death. And in recent years, both unintentional and intentional self-poisoning admissions per million packs of paracetamol sold have been decreasing.

## How might paracetamol access change?

The [expert panel](#) made several [recommendations](#) including:

- limiting pack sizes
- introducing purchase limits
- requiring a prescription to purchase higher amounts and modified-release products (such as Panadol Osteo)
- requiring a prescription for people under 18.

[Additional measures](#) being considered include changes to packaging and how paracetamol is displayed in stores. At the moment, there are no recommendations to make all paracetamol products prescription-only, or to restrict their sale to pharmacies only.

Many of these strategies are focused on reducing potential harm by limiting the amount of paracetamol available. The change to pack sizes available outside of pharmacies would bring Australia in line with countries such as Denmark, Ireland and the United Kingdom. These countries have had restricted pack sizes (maximum 10–16 tablets) for sale outside pharmacies for many years and seen [reductions](#) in [hospital admissions](#) and [deaths](#). Poisonings among people aged 10–17 years [also went down](#) in Denmark after age restrictions were introduced in 2011.

The [expert report found](#) most cases of self-poisoning involve medicines already present at home. So the benefits of these measures may be limited. But limiting pack sizes may help to reduce the overall quantity of medicines available in the home and the risk of a fatal poisoning.

The expert panel also recommended better follow-up care after self-poisoning events. Developing [preventative strategies](#) and [increasing mental health support](#) is vital to address the drivers of intentional self-poisonings more broadly.

## What next?

Ready access to paracetamol allows people to conveniently self-treat

their pain at minimal cost and without visiting a health professional. Reduced paracetamol availability may prompt people to switch to using other pain medicines, such as ibuprofen, which [carries its own risks](#) and may not be suitable for everyone.

Although limiting harms due to paracetamol poisoning is important, at the same time, the TGA will need to ensure paracetamol remains accessible to those in need. By keeping paracetamol on the shelves of supermarkets and pharmacies in reduced pack sizes, the TGA will be aiming to strike a balance between accessibility and safety.

The biggest potential impact for people living with chronic pain would be brought about by reducing larger pack sizes currently only available in pharmacies (such as those containing 50–100 tablets) and making modified-release paracetamol products prescription-only. The [expert report](#) notes that at 665 mg, the modified-release formulation in particular is linked to larger overdoses. As these products are currently only available from pharmacies, authorities would need to show how these changes would significantly reduce harm, without overly burdening people in pain.

The [TGA is seeking feedback](#) until mid-October to guide its decision. After that the TGA's expert advisory committee will consider whether to amend the Poisons Standard to change how paracetamol can be accessed. For now, paracetamol is available as usual.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Restrictions on paracetamol may be coming. What does that mean for consumers?



(2022, September 29) retrieved 26 April 2024 from  
<https://medicalxpress.com/news/2022-09-restrictions-paracetamol-consumers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.