


# **Schizophrenia and bipolar disorder may be detectable years before illnesses begin**

September 8 2022

---



Credit: Pixabay/CC0 Public Domain

The risk of schizophrenia and bipolar disorder may be detectable years before the illnesses begin, according to new research.

A University College Dublin led study funded by the Health Research Board has found that 50% of people who developed these [mental health disorders](#) had attended specialist child and adolescent mental health services (CAMHS) in [childhood](#).

Published in the journal *World Psychiatry*, the findings suggest the possibility of earlier intervention and even prevention according to Professor Ian Kelleher, from the UCD School of Medicine, who led the international study which was carried out in conjunction with the Finnish Institute for Health and Welfare (THL).

"Schizophrenia and bipolar disorder typically emerge in early adulthood and can have a devastating impact on the [individuals](#) affected, as well as on their families," he said.

"Our findings show that half of individuals who develop these illnesses had come to CAMHS at some stage in childhood, typically many years before they developed schizophrenia or bipolar disorder.

"We know that early intervention is key to improving outcomes for people with [serious mental illness](#). These findings demonstrate the enormous opportunities to provide far earlier intervention, even while still in childhood, by developing specialist early intervention services within existing child and adolescent mental health services".

Schizophrenia and bipolar disorder are serious mental illnesses affecting



about 65 million people worldwide. Both disorders are usually diagnosed in adulthood and are often associated with high levels of disability, personal and societal cost. Early intervention, however, is known to lead to better outcomes for people affected by these illnesses.

The researchers behind the new study used Finland's world-leading healthcare registers to trace all individuals born in 1987 throughout childhood and adolescence to see if, between birth and age 17 years, they ever attended CAMHS.

Using unique patient identifiers, the researchers were then able to follow all these individuals up to age 28 years and see who went on to be diagnosed with schizophrenia or bipolar disorder.

They found that the risk of psychosis or [bipolar disorder](#) by age 28-years-old was 1.8% for individuals who had not attended CAMHS. For individuals who had attended outpatient CAMHS in adolescence, however, the risk was 15% and for individuals who had been admitted to an inpatient adolescent CAMHS hospital, the risk was 37%.

"This research shows the power of electronic healthcare registers to answer important questions about [human health](#) and disease," said Professor Mika Gissler, THL.

"It demonstrates how healthcare register data can be used to better understand pathways to serious mental illness, from childhood into adulthood, and to identify critical opportunities for early [intervention](#)."

Stressing the importance of [early intervention](#), Professor Ian Kelleher said: "We know it's crucial to intervene as early as possible to prevent some of the worst effects of these illnesses. But ideally, we would like to be able to intervene even before the onset of [illness](#), to prevent it altogether.

These findings highlight the possibility of intervening far earlier than we do at present, even in childhood and adolescence, to prevent these serious mental illnesses from emerging".

**More information:** Potential for prediction of psychosis and bipolar disorder in Child and Adolescent Mental Health Services: a longitudinal register study of all people born in Finland in 1987, *World Psychiatry* (2022). [osf.io/gku9a/](https://osf.io/gku9a/)

Provided by University College Dublin

Citation: Schizophrenia and bipolar disorder may be detectable years before illnesses begin (2022, September 8) retrieved 7 May 2024 from <https://medicalxpress.com/news/2022-09-schizophrenia-bipolar-disorder-years-illnesses.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--