


Scientists find cold spells in the tropics increase heart attack risk

September 2 2022



Graphical abstract. Credit: *Science of The Total Environment* (2022). DOI: 10.1016/j.scitotenv.2022.158010

Scientists have long known that temperature changes impact the incidence of heart attacks. However, most of the research done so far has been in temperate climates, where temperatures range widely.

Now, a team of researchers has investigated how the narrow temperature ranges of a tropical climate impact the incidence of a specific type of myocardial infarction (the medical term for a [heart attack](#)) in Singapore. The researchers say their findings, published in the journal *Science of the Total Environment*, could have health policy implications for populations within cosmopolitan cities in the tropics.

"Using 10 years of nationally collected data, we found strong evidence that a drop of 1°C in [ambient temperature](#) increased the risk of a type of acute myocardial infarction in the population by 12%," said co-senior author Professor Marcus Ong, Director of the Health Services & Systems Research Program and the Pre-hospital & Emergency Research Center (PERC) at Duke-NUS Medical School.

"Furthermore, people aged 65 and above appeared to be about 20% more vulnerable to cooler temperatures compared to [younger people](#)," added Prof. Ong, who is also Senior Consultant at the Department of Emergency Medicine at Singapore General Hospital (SGH).

The study, which was conducted in collaboration with Singapore's National Environment Agency (NEA), analyzed daily patient records from the Singapore Myocardial Infarction Registry. The researchers were specifically looking for people who experienced non-ST-segment elevation myocardial infarction (NSTEMI). This is a type of acute heart attack that happens when a blood vessel feeding the heart becomes partially blocked. When doctors examine the patient's electrocardiogram (ECG) results, they don't find the easily identifiable ST elevation that signifies another type of heart attack, STEMI, which occurs when the coronary artery is completely blocked. Since the 1980s, the incidence of

NSTEMI has risen while that of STEMI has declined.

The researchers were able to collect 60,643 reports of NSTEMI between 2009 and 2018. They then statistically analyzed how the onset of NSTEMI in these patients correlated with local meteorological data obtained from [weather stations](#) across Singapore, including mean temperature and rainfall.

Cooler ambient temperatures were independently associated with an increased risk of NSTEMI up to 10 days after a temperature drop. There were no [gender differences](#) relating to the effects of warmer or cooler temperatures on NSTEMI risk. Nor were changes in rainfall associated with an increased risk.

"Our study found that even in a relatively warm part of the world, cooler ambient temperatures increased the risk of heart attacks," said Dr. Andrew Ho, one of the study's first authors, who is an Assistant Professor with PERC and an Associate Consultant with SGH's Department of Emergency Medicine. "This improves our understanding that deviations from the temperature that one is used to can lead to harmful bodily stress. Consistent with our previous studies that showed that the elderly were more susceptible to environmental stressors including air pollution, we found some evidence that this group of individuals were at greater risk of heart attacks at [cooler temperatures](#)."

"There are several individual-level risk factors for cardiovascular disease, but none are as widely experienced as [weather patterns](#)," said Dr. Joel Aik, an environmental epidemiologist and co-senior author of the study from NEA, who is also an Adjunct Assistant Professor with PERC. "Daily weather variations have the capacity to trigger [cardiovascular disease](#) events in at-risk individuals, with particular implications for Singapore's aging population. In the context of climate change, these findings highlight a risk factor of substantial public health

concern."

Further research over a longer period is needed to confirm the results. The team also recommends research that helps to identify the biological pathways involved in increased vulnerability of the elderly to cold-related NSTEMI in the tropics.

More information: Annabel Seah et al, Ambient temperature and hospital admissions for non-ST segment elevation myocardial infarction in the tropics, *Science of The Total Environment* (2022). [DOI: 10.1016/j.scitotenv.2022.158010](https://doi.org/10.1016/j.scitotenv.2022.158010)

Provided by Duke-NUS Medical School

Citation: Scientists find cold spells in the tropics increase heart attack risk (2022, September 2) retrieved 30 June 2024 from <https://medicalxpress.com/news/2022-09-scientists-cold-tropics-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.