

Vaccinating women infected with COVID during pregnancy prior to delivery provides antibodies to newborns

September 26 2022



Credit: Unsplash/CC0 Public Domain

Women with COVID in pregnancy who are subsequently vaccinated after recovery, but prior to delivery, are more likely to pass antibodies on

to the child than similarly infected but unvaccinated mothers are. Researchers who studied a mix of vaccinated and unvaccinated mothers found that 78% of their infants tested at birth had antibodies. Of these infants, 3 of 4 born to unvaccinated mothers had evidence of antibodies while all of those from vaccinated mothers carried COVID antibodies.

At six months, 52% of the babies, from both vaccinated and unvaccinated mothers, carried antibodies. The decrease in that percentage is due to a combination of infants of vaccinated mothers carrying more antibodies than those from unvaccinated mothers at [birth](#), a waning of detectable antibodies over time, and some participants leaving the trial prior to the 6-month point. Some infants born to unvaccinated mothers had no detectable antibodies at birth.

There are currently no approved COVID vaccines for newborns under the age of six months. Transplacental transfer of antibodies to the infant from the mother during pregnancy may provide protection against COVID during the first six months of life. Also, little is known about the impact of mothers' COVID severity, timing of infection and subsequent vaccination on both maternal and infant antibodies over time.

In a [longitudinal study](#), researchers collected [blood samples](#) from [pregnant women](#) and infants at the time closest to infection, birth, and six months postpartum: samples were collected from 148 women and 122 newborns at birth, and another 45 maternal and 48 infant samples were taken at six months. Some participants dropped out of the trial during the intervening time, and some mothers gave birth to twins or triplets, accounting for the differences in numbers tested.

This is one of the largest longitudinal studies of mothers and infants with a history of COVID infection during the mother's pregnancy. However, the researchers note some limitations to the study. Due to the study's design, associations may not imply causation, they did not have

vaccinated controls without a history of COVID infection for comparison, and they had a high attrition rate in study subjects by the 6-month point.

In this analysis, the mother's vaccination status was the strongest predictor of antibody transfer to the infants, who were more likely to have detectable antibodies at birth. This may be an [effective strategy](#) to boost COVID antibodies not just in mothers, but in [infants](#) before they are eligible for the vaccine at six months of age.

"Several studies have demonstrated that mothers with a history of COVID during pregnancy may pass antibodies to their babies at delivery," said lead author Dr. Mary Cambou, clinical instructor in the UCLA Division of Infectious Diseases. "However, our study was able to examine several factors, including timing, the severity of COVID disease, and subsequent vaccination, on both maternal and infant levels at birth and at six months. Vaccination following infection and prior to delivery was the strongest predictor of infant [antibodies](#) at birth. Notably, infant levels dropped significantly by six months, emphasizing the importance of starting the COVID vaccine series as early as six months."

The study is published in the *Journal of Infectious Diseases*.

More information: Mary C Cambou et al, Longitudinal Evaluation of Antibody Persistence in Mother-Infant Dyads Following SARS-CoV-2 Infection in Pregnancy, *The Journal of Infectious Diseases* (2022). [DOI: 10.1093/infdis/jiac366](https://doi.org/10.1093/infdis/jiac366)

Provided by University of California, Los Angeles

Citation: Vaccinating women infected with COVID during pregnancy prior to delivery provides antibodies to newborns (2022, September 26) retrieved 3 May 2024 from <https://medicalxpress.com/news/2022-09-vaccinating-women-infected-covid-pregnancy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.