


Advancing the acute treatment of stroke

October 13 2022



Distribution of functional outcomes at 90 days in the intention-to-treat population. Credit: *New England Journal of Medicine* (2022). DOI: 10.1056/NEJMoa2206317

Stroke is the "number one killer" that endangers life and health in China. Statistics released by the National Health Commission show that the total number of stroke patients in China exceeds 28 million, and at least 1 in every 5 deaths is due to stroke.

The latest "China Stroke Report 2020" indicates that there are about 3.94 million new strokes and 2.19 million deaths due to [stroke](#) every year in China. It is estimated that the number of stroke patients in the

world has exceeded 100 million. The World Stroke Organization has designated October 29 as "World Stroke Day".

Early re-canalization of occluded vessels following acute ischemia stroke is key to improving prognosis. At present, the most effective treatment methods for [acute ischemic stroke](#) are intravenous thrombolysis and intra-arterial [thrombectomy](#).

A new study, led by Prof. Wei Hu, director of the stroke center and department of neurology at the first affiliated hospital of the University of Science and Technology of China has demonstrated that in patients who suffered a basilar arterial [occlusion](#), an endovascular thrombectomy in 12 hours will lead to a better functional outcome compared with standard medical therapy.

In addition, endovascular thrombectomy can significantly improve patients' self-care ability and reduce mortality. This study is the first clinical trial successfully demonstrating the superiority of endovascular thrombectomy in acute basilar artery occlusion.

The research team has named the clinical trial "ATTENTION" in the hope that prevention, treatment and research of strokes will attract an increasing amount of attention. Among all types of stroke, posterior circulation stroke accounts for about one fifth. Acute ischemic stroke caused by basilar artery occlusion is often associated with a poorer prognosis and a higher mortality than intracranial large vessel occlusion in the anterior circulation.

However, due to the lack of high-quality evidence-based [clinical trials](#), the efficacy and safety of endovascular thrombectomy for acute basilar artery occlusion has been difficult to determine. Before the launch of this study, the treatment team of the neurology department of the First Affiliated Hospital of the University of Science and Technology of

China had conducted a four-year clinical observational registry trial. The initiation of ATTENTION trial was based on high-quality evidence from the results of the preceding observational trial.

The research was published in the *New England Journal of Medicine*.

More information: Chunrong Tao et al, Trial of Endovascular Treatment of Acute Basilar-Artery Occlusion, *New England Journal of Medicine* (2022). [DOI: 10.1056/NEJMoa2206317](https://doi.org/10.1056/NEJMoa2206317)

Provided by University of Science and Technology of China

Citation: Advancing the acute treatment of stroke (2022, October 13) retrieved 7 May 2024 from <https://medicalxpress.com/news/2022-10-advancing-acute-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--