


Assumptions about the lethality of air pollution in India may be exaggerated

October 12 2022



Credit: *Environmental Health Perspectives* (2022). DOI: 10.1289/EHP9538

High levels of exposure to air pollution in India have a smaller effect on

mortality than previously estimated, according to the first study to cover the whole of India. The research was recently published in *Environmental Health Perspectives*.

India has among the highest levels of [air pollution](#) in the world, and nearly every Indian lives in areas with [fine particulate matter](#) levels well above PM2.5—the level considered safe by the World Health Organization (WHO).

The nationwide study linked concentrations of PM2.5 derived from satellite-based measurements of deaths in over 7,400 small areas among seven million people.

The authors quantified the relationship between PM2.5 exposures over several years to the subsequent mortality risk, focusing on the leading killers of Indian adults: respiratory disease, [heart disease](#), stroke and overall mortality. Their analyses adjusted for risk factors that make these diseases more common such as smoking, urban or rural residency, and education. Because these diseases and PM2.5 tend to cluster in some areas but not others, they also adjusted for spatial clustering.

Analyzing over 200,000 deaths at ages 15–69 years, the authors found a nine percent excess risk in stroke deaths for every 10-unit (measured as micrograms per cubic meter) increase in PM2.5 exposure. However, there were no excess risks for respiratory or heart disease and for total mortality (after excluding stroke). Study results were broadly similar in different age groups and in areas with higher or lower use of household solid fuel (which itself is the major contributor to PM2.5 exposure as measured by satellites).

The MDS study results showing little or no effect on heart disease and only modest effects on stroke were similar to an earlier smaller [study](#) led by McMaster University in Canada that examined air pollution and

deaths in 750 communities in 21 countries. In contrast to these [direct observations](#), earlier estimates of air pollution deaths in India use complex models that assume a much higher mortality risks from PM2.5 exposure.

"Measuring the health effects of air pollution is complicated," says lead author Prof. Patrick Brown, from the University of Toronto's Department of Statistics. "There are so many things to tease out to establish if a relationship with deaths is true, or just noise in the data. Even the modest excess risks we found for stroke might reflect the uncertainty that is inherent in these types of epidemiological studies."

Brown added, "Direct studies like the MDS are far preferable to extrapolating from models, many of which take reports from non-Indian settings and apply them—unseeingly—to India. Earlier models have probably overstated the extent to which air pollution kills adults in India."

"Most existing studies on air quality and health are done in high-income countries, where PM2.5 exposures are far lower and disease patterns are substantially different than in low and middle-income countries such as India, says co-author Prof. George D'Souza, Dean of St. John's Medical College, Bengaluru, India. "That needs to change."

The results are one of many insights being generated by the [Million Death Study](#)—one of the largest studies of premature mortality in the world. The study, covering deaths from 2001 to 2014, is a long-running effort to better understand the causes of mortality in countries such as India, where the vast majority of people die at home without a [death certificate](#). Launched by Jha at DLSPH, it has helped governments and health providers to better allocate scarce public health funding.

Despite the good news, there are many reasons to act on air pollution in

India. "PM2.5 may not be the big killer in India as has been touted, but action to reduce PM2.5 exposure is still necessary," says Jha, who is also a physician at St. Michael's Hospital, a site of Unity Health Toronto. "Air pollution worsens child lung health and lowers quality of life, and that is sufficient justification to act."

More information: Patrick E. Brown et al, Mortality Associated with Ambient PM2.5 Exposure in India: Results from the Million Death Study, *Environmental Health Perspectives* (2022). [DOI: 10.1289/EHP9538](https://doi.org/10.1289/EHP9538)

Provided by University of Toronto

Citation: Assumptions about the lethality of air pollution in India may be exaggerated (2022, October 12) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-10-assumptions-lethality-air-pollution-india.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.