

# Binge eating is more common than anorexia or bulimia—but it remains a hidden and hard-to-treat disorder

October 23 2022, by Hannah Kennedy

---



Credit: Unsplash/CC0 Public Domain

For many people, the term "eating disorder" will bring to mind its two most familiar forms—anorexia nervosa and bulimia nervosa. What they

may not realise, however, is that "binge eating disorder" is [more common than the other two combined](#) and can significantly reduce quality of life.

More than just eating too much, which most people will find themselves doing once in a while, binge eating reflects a recurrent behaviour pattern of losing control, bingeing food and feeling associated shame and guilt afterwards.

Binge eating disorder was only recognised as a diagnosis in [2013](#), much later than anorexia and bulimia. But a lack of awareness about binge eating means people who experience the disorder aren't getting the help they need from doctors.

As our [multinational research project](#) has recently found, people with binge eating disorder are also underrepresented in studies of eating disorders, limiting the ability of researchers to develop treatments.

## Understanding the fundamentals of binge eating

Unlike bulimia, or the type of anorexia that involves binge eating and purging, people experiencing binge eating disorder don't try to compensate with strategies such as purging or excessive exercise following a bingeing episode.

Also unlike anorexia or bulimia, which predominantly affect woman, binge eating disorder affects an estimated [3.5% of women and 2% of men](#) during their lifetimes. In New Zealand, that could mean more than 130,000 people suffer from this particular eating disorder. In Australia, this could affect more than 600,000 people in their lifetimes.

Although it may first emerge in childhood and adolescence, binge eating more typically occurs in [early adulthood](#). Importantly, bingeing can be

observed in people of all body sizes, although many individuals who experience binge eating will be in a [higher weight range](#).

According to the US [National Library of Medicine](#), there are five criteria to diagnose binge eating disorder:

1. Recurrent episodes of binge eating characterised by eating, in a limited period of time, a larger amount of food than most people would eat under similar circumstances, while also experiencing a lack of control over eating during the episode.
2. Binge eating episodes are associated with three or more of the following factors: eating much more rapidly than normal, eating until feeling uncomfortably full, eating large amounts of food when not feeling physically hungry, eating alone due to embarrassment and feeling disgusted, depressed or very guilty afterwards.
3. Marked distress caused by binge eating.
4. Binge eating, on average, at least once a week for three months.
5. The binge eating is separate from other disordered eating behaviours, including vomiting after eating, as happens with bulimia.

Other warning signs of binge eating disorder include:

- frequent weight fluctuations
- fad diets, including eliminating entire food groups (carbs, sugar, dairy, etc)
- extreme concern with [body weight](#) and shape
- stealing or hoarding food
- withdrawal from friends and usual activities
- eating in secret and hiding evidence (such as food wrappers).

The longer-term [health complications](#) associated with this type of eating

disorder include heart disease, type 2 diabetes, obesity and osteoarthritis (due to wearing down of the cartilage that cushions the ends of bones at the joints).

## Hidden from view

Despite all this, binge eating is still considered the "hidden" eating disorder. Without the extreme weight loss or purging behaviours that can accompany other eating disorders, binge eating is less recognisable.

Bingeing may even be dismissed as just an occasional over-indulgence. Media portrayals of eating disorders perpetuate this idea, often focusing on anorexia.

The stigma and shame surrounding binge eating behaviours are significant and may prevent people from accessing help and treatments that could combat the disorder. Fewer than half of those who experience binge eating will [seek and receive treatment](#).

Patients will often not disclose their binge eating symptoms to their doctor, meaning treatments often focus on weight loss and [other health complications of binge eating](#) but fail to recognise the eating disorder that drives the behaviour.

This lack of awareness and discussion affects recruitment into research in eating disorders.

## The research gap

Recruiting people who have experienced binge eating for eating disorder studies is critical for better understanding of the disorder and improving treatments.

However, our experience of recruiting participants for our own study indicates this can be challenging. Despite our best efforts, the number of participants with anorexia or bulimia far outnumber those with binge eating—a pattern observed in New Zealand, Australia and the US.

[Current treatments](#) for adults with binge eating disorder include [cognitive behavioural therapy](#), antidepressants and, in some countries, the drug [lisdexamfetamine](#), which is the only approved medication for the disorder so far.

## Visibility will improve treatment options

Awareness and education that binge eating is a prevalent—and potentially life-threatening eating disorder—is desperately needed to combat the current stigma and provide insight into the drivers of the behaviour.

The government announced an additional [NZ\\$3.9 million](#) in funding for eating disorder services in this year's budget. But if binge eating remains hidden from view, sufferers could miss out on the essential support this boost could provide.

People with [binge eating](#) disorder should be supported by their doctors to seek help and encouraged to engage in research that will eventually lead to better outcomes. Until that happens, we are working blind as we try to address a disorder that affects thousands.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Binge eating is more common than anorexia or bulimia—but it remains a hidden and hard-to-treat disorder (2022, October 23) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-10-binge-common-anorexia-bulimiabut-hidden.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.