

Black patients with atrial fibrillation more likely to experience adverse outcomes

October 27 2022



Credit: Pixabay/CC0 Public Domain

Black patients with atrial fibrillation are less likely to be prescribed newer anticoagulant medications when they leave the hospital compared to their white counterparts and are consequently more likely to

experience adverse outcomes, including stroke and death, according to a new study led by University of Pittsburgh School of Medicine physician-scientists and published today in *JAMA Cardiology*.

This research builds on a study published in 2018 that showed Black and Hispanic patients were less likely to receive newer [anticoagulant](#) medications than [white patients](#). In the new study, researchers directly correlate this disparity in prescribing to [adverse outcomes](#) in Black and Hispanic populations.

"We knew that Black patients were less likely to receive anticoagulants, but our goal was to determine how this difference in treatment affected [patient outcomes](#)," said Utibe Essien, M.D., M.P.H., assistant professor of medicine at Pitt. "With access to such a large registry, we were able to retrospectively look at patients from across the nation at different types of hospitals to make that link between treatment and outcomes. The results are truly concerning."

Atrial fibrillation is the most common type of heart arrhythmia, contributing to approximately 160,000 deaths in the United States each year. Patients with atrial fibrillation are at a higher risk for stroke and heart failure. The number of patients with atrial fibrillation continues to rise, and an estimated 12.1 million people in the United States will have the condition by 2030.

The study used the "Get With the Guidelines" registry to look at nearly 70,000 patients who had been hospitalized and diagnosed with [atrial fibrillation](#) between 2014 and 2020. Consistent with previous research, the study found that Black patients were 25% less likely to receive an anticoagulation medication at discharge compared to white patients, and they were even less likely to receive newer and more effective oral anticoagulants. In a follow-up a year later with 16,000 of these original patients, the researchers showed that Black patients had double the risk

of a stroke, as well as a 1.2 times higher risk of death. Similar [stroke](#) results were found for Hispanic patients.

Determining why these disparities exist among Black and Hispanic patients is a more complex question. While researchers found that the size or location of the hospital where the patient was admitted did not have an effect on patient outcomes, access to high-quality cardiovascular care may play a part.

However, even if patients have access to cardiovascular care, social and economic influences play a major role in determining if patients can receive medications. Direct anticoagulants, while newer and more effective, can often cost patients \$500 a month, making them difficult for lower-income patients to afford.

In addition, the trust between a patient and provider can impact whether patients are prescribed medications and if they take them. While [implicit bias](#) may be a factor in whether patients are prescribed newer medications, if patients do not trust their provider, they may be less likely to take the anticoagulants even if they were prescribed.

"These results give us an opportunity to think deeply about pharmaco-equity and the steps we can take urgently to improve treatment and outcomes for patients," Essien said. "We need to continue to prioritize policies that make it easier for more patients to afford the [medication](#) they need, and work with providers to ensure we're treating patients in a way that builds trust and leads to better outcomes."

More information: Utibe R. Essien et al, Association of Race and Ethnicity With Oral Anticoagulation and Associated Outcomes in Patients With Atrial Fibrillation, *JAMA Cardiology* (2022). [DOI: 10.1001/jamacardio.2022.3704](https://doi.org/10.1001/jamacardio.2022.3704)

Provided by University of Pittsburgh Schools of the Health Sciences

Citation: Black patients with atrial fibrillation more likely to experience adverse outcomes (2022, October 27) retrieved 20 June 2024 from <https://medicalxpress.com/news/2022-10-black-patients-atrial-fibrillation-adverse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.