

Broken bones, fractured skulls remain common bicycle injuries, especially among teen boys: 20-year review

October 7 2022



Credit: CC0 Public Domain

More than one million U.S. children fractured a bone while riding a bicycle over the past 20 years, and most of them were boys between ages 10 to 15 years old, according to research presented during the 2022

American Academy of Pediatrics National Conference & Exhibition.

Authors of the abstract, "Pediatric Fractures Associated with Riding Bicycles: A National Twenty-Year Analysis," found that, despite efforts to make roads more bike-friendly, over 65,000 injuries occurred in an accident involving a motor vehicle. About 85% of children who sustained a skull fracture while riding bicycles were not wearing helmets.

"The results of our study suggest that continued efforts teaching road safety and promoting helmet use should be targeted towards all children, but with additional efforts being directed towards the most affected population, namely 10- to 15-year-old boys," said J. Todd R. Lawrence, MD, Ph.D., at Children's Hospital of Philadelphia. "Municipalities should continue to evaluate traffic patterns on their local roads to improve bike safety for children."

The researchers examined information provided in the National Electronic Injury Surveillance System database for the years 2001 through 2020 to identify patients ages 18 or younger who showed up at U.S. emergency departments with fractures associated with bicycles. They found an average 50,975 fractures reported annually, with about 71% of patients being male. Only a small minority of patients with [skull fractures](#) were wearing helmets; nearly 87% of patients were not wearing [helmets](#) at the time of the skull fracture.

While the number of fractures from children riding bicycles has decreased over the past two decades, researchers noted an increase in fractures in 2020 that is consistent with literature showing a significant increase in other injuries during the COVID-19 pandemic. This is potentially due to an increased use of [bicycles](#) from stay-at-home orders and cancellation of school and summer camps.

"Given the results of our study, we recommend targeting bicycle safety

efforts toward the most affected populations, largely 10- to 15-year-old boys," said William Huffman, a medical student. "Teaching road and helmet safety for [bicycle](#) riders is paramount to keeping children safe."

More information: Pediatric Fractures Associated with Riding Bicycles: A National Twenty-Year Analysis, 2022 American Academy of Pediatrics National Conference & Exhibition

Provided by American Academy of Pediatrics

Citation: Broken bones, fractured skulls remain common bicycle injuries, especially among teen boys: 20-year review (2022, October 7) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-10-broken-bones-fractured-skulls-common.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--