

# Study explores links between people taking multiple medications and dementia diagnosis

October 12 2022

---



Credit: CC0 Public Domain

People with dementia are likely to have taken more than three medications for other health conditions in the five years directly before their diagnosis, according to new research.

The study is the first to provide an in-depth exploration of the links

between evolving polypharmacy—which involves a patient being prescribed more than one drug at any given time—and a dementia diagnosis.

Published in the *Aging and Disease* journal, it is based on an analysis of the records of more than 33,000 [dementia patients](#) in Wales between 1990 to 2015.

Experts in e-health used machine learning techniques to identify potentially damaging patterns in a patient's medicine usage, and how these patterns evolve in the run-up to diagnosis.

They found that in the 20 years leading up to them being diagnosed, the proportion of patients taking three or more medications rose from 5.5% (for the period 16 to 20 years prior to diagnosis) to 82.16% among those less than five years from a diagnosis.

Researchers also found that as the development towards dementia progressed, the patterns of polypharmacy shifted from being clearly distinct to being more closely associated with particular medical conditions.

And of those closest to their [diagnosis](#), almost two-thirds (66.55%) were found to be taking multiple medicines for a combination of respiratory or urinary infections, arthropathies and rheumatism, and cardio-vascular disease. A further 22% of patients were found to be taking medicines for infections, arthropathies and rheumatism, cardio-metabolic disease and depression.

The study was supported by the Health Data Research UK and conducted by an international team of researchers from the University of Plymouth, Aptuit (an Evotec company), Swansea University Medical School, and the University of Oxford.

Shangming Zhou, Professor of e-Health at the University of Plymouth, led the study. He said: "Given the rise in dementia cases internationally, the need to understand how patterns of polypharmacy evolve before and after a [dementia diagnosis](#) are important for devising a safe treatment program for each patient. Our aim in this study was to help doctors find ways to prescribe multiple items of dementia [medication](#) safely and without reducing their effectiveness. The use of machine learning has been vital in helping us understand how these patterns develop, and our hope is we can now use this knowledge to treat patients."

It has previously been established that when multiple types of preventative medication are being prescribed, the benefits of the drugs may be reduced and the chances of harm from drug interaction and side effects increased.

Those requiring hospital treatment who are taking multiple medications are also known to have a higher likelihood of re-admission within three months after being discharged.

With the number of people with [dementia](#) in the UK projected to rise to 1.6 million by 2040, researchers hope this new study will inform safe prescribing practices and encourage doctors to prescribe medicine combinations developed with a view to minimizing cognitive impairments.

**More information:** Identifying Dynamic Patterns of Polypharmacy for Patients with Dementia from Primary Care Electronic Health Records: A Machine Learning Driven Longitudinal Study, *Aging and Disease* (2022). [DOI: 10.14336/AD.2022.0829](https://doi.org/10.14336/AD.2022.0829)

Provided by University of Plymouth

Citation: Study explores links between people taking multiple medications and dementia diagnosis (2022, October 12) retrieved 4 May 2024 from <https://medicalxpress.com/news/2022-10-explores-links-people-multiple-medications.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.