

'Front-loading' alcohol, an important focus for future research

October 14 2022



Credit: Unsplash/CC0 Public Domain

Front-loading alcohol, or drinking faster at the outset of a drinking session, may be associated with harmful patterns which can lead to alcohol use disorders and should be studied, according to the authors of

an analysis published in *Alcoholism: Clinical & Experimental Research*. The analysis suggests increasing evidence that front-loading drinking behavior is driven by reward effects more than by other factors, such as metabolic tolerance or negative reinforcement.

Some studies have suggested that the rate of drinking, not just the quantity of alcohol consumed, is correlated with symptoms of alcohol use disorder. Front-loading is not uniformly defined in the literature and is sometimes conflated with other drinking behaviors, such as binge drinking and 'pre-gaming' or 'pre-partying.' However, these behaviors don't necessarily meet the criteria for front-loading, which the authors define as a pattern of [behavior](#) where the subject drinks more during the beginning of access to alcohol and consumes enough alcohol during that period to quickly produce a pharmacological effect, such as a 'buzz.'

The analysis reviewed studies that contain reports of front-loading behavior in animal or [human subjects](#) to determine possible motivations for front-loading behavior. Limited but increasing evidence suggests that the rewarding effects of alcohol drive front-loading behavior, where individuals learn that quicker consumption of alcohol leads to more rapid intoxication and, usually, greater reward. There were insufficient data to identify negative reinforcement, that is, [drinking](#) to relieve anxiety, stress, or [withdrawal symptoms](#), as a motivation for front-loading. However, more research is needed.

The authors did not find evidence to support metabolic tolerance to alcohol as a driver of front-loading but suggest that chronic behavioral tolerance, where the brain learns to counteract the effects of alcohol, cannot be ruled out. Drinking faster may allow individuals to, for a time, overcome that tolerance and more strongly feel alcohol's desired effects. Studies to effectively assess behavioral tolerance are recommended.

Front-loading is not well studied but may be important for understanding

the causes of [alcohol](#) use disorder. The authors recommend research to address knowledge gaps related to front-loading, including [gender differences](#) in front-loading, different contexts for front-loading, acute tolerance, and negative reinforcement as motivations for front-loading, and better assessment of behavioral tolerance.

More information: Cherish E. Arding et al, A critical review of front-loading: A maladaptive drinking pattern driven by alcohol's rewarding effects, *Alcoholism: Clinical and Experimental Research* (2022). [DOI: 10.1111/acer.14924](https://doi.org/10.1111/acer.14924)

Provided by Research Society on Alcoholism

Citation: 'Front-loading' alcohol, an important focus for future research (2022, October 14) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-10-front-loading-alcohol-important-focus-future.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--