

Likelihood of receiving an autism diagnosis may depend on where you live

October 24 2022



Credit: Pixabay/CC0 Public Domain

New autism diagnoses in the UK tend to be clustered within specific NHS service regions, suggesting that where an individual lives may influence whether they receive an autism diagnosis and access to special

education needs support.

The latest findings, from researchers from the University of Cambridge in collaboration with researchers from the London School of Economics and Political Science and Newcastle University, are published today in *The Lancet Child & Adolescent Health*.

After analyzing all new autism cases across England using NHS health service boundaries for possible hotspots, some areas stand out. For example, 45.5% of the NHS Rotherham catchment area had higher-than-average new autism diagnoses clusters. For NHS Heywood, this amounted to 38.8% of its catchment area and 36.9% for NHS Liverpool, pointing at a possible health service effect towards who receives an [autism diagnosis](#).

The research team used four years' worth of data from the Summer School Census, which collected data from individuals aged 1-18 years old in state-funded schools in England. Of the 32 million pupils studied, more than 102,000 new autism diagnoses were identified between 2014 and 2017.

After adjusting for age and sex, the researchers found that one in 234 children were given a new autism diagnosis during that four-year period. New diagnoses tended to happen when children are transitioning to a new school, whether that was into nursery (1-3 years), primary school (4-6), and secondary school (10-12 years).

Particular communities appeared to have different rates, varying by ethnicity and deprivation.

Lead researcher Dr. Andres Roman-Urrestarazu from the Department of Psychiatry and Cambridge Public Health at the University of Cambridge said, "Autism diagnoses are more common among Black students and

other minority [ethnic groups](#). Why this is the case is not clear and so we need to explore the role played by [social factors](#) such as ethnicity and area deprivation as well as the nature of local services."

The likelihood of receiving an autism diagnosis more than tripled among girls depending on their ethnicity and social and financial situation, compared to white girls without financial disadvantages who speak English as their first language.

In contrast, boys' likelihood of receiving an autism diagnosis increased more than five-fold depending on their ethnicity and social and [financial situation](#), compared to white boys without financial disadvantages who speak English as their first language.

Boys and young men are already known to be more likely to receive autism diagnoses, but the social determinants that could affect a diagnosis remained an open question.

Dr. Robin van Kessel, co-lead researcher from the Department of Health Policy at the London School of Economics and Political Science, said, "These new findings show how [social determinants](#) interact and can combine to significantly increase the likelihood of an autism diagnosis. As a result, individuals from a minority ethnic background experiencing [economic hardship](#) may be significantly more likely to receive an autism diagnosis than their peers."

Professor Carol Brayne from Cambridge Public Health said, "There are clear inequalities in an individual's likelihood of receiving an autism diagnosis, whether they are socioeconomic factors, ethnicity or even which NHS region or local authority someone lives in."

More information: Autism incidence and spatial analysis in more than 7 million pupils in English schools: a retrospective, longitudinal, school

registry study, *The Lancet Child & Adolescent Health* (2022). [DOI: 10.1016/S2352-4642\(22\)00247-4](https://doi.org/10.1016/S2352-4642(22)00247-4)

Provided by University of Cambridge

Citation: Likelihood of receiving an autism diagnosis may depend on where you live (2022, October 24) retrieved 3 May 2024 from <https://medicalxpress.com/news/2022-10-likelihood-autism-diagnosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.