

## **Engineers record neurons to pinpoint synaptic links**

October 19 2022, by Jade Boyd



Using millisecond-scale recordings from a dense 3D array of electrodes, researchers in the laboratory of Rice University neuroengineer Chong Xie were able to map the locations and activity of up to 1 million potential synaptic links in a living brain. Credit: Xie Lab/Rice University



It's a mystery how human thoughts and dreams emerge from electrical pulses in the brain's estimated 100 trillion synapses, and Rice University neuroengineer Chong Xie dreams of changing that by creating a system that can record all the electrical activity in a living brain.

In a recently published study in *Nature Biomedical Engineering*, Xie and colleagues described their latest achievement toward that goal, a 3D electrode array that allows them to map the locations and activity of up to 1 million potential synaptic links in a living brain based on recordings of the millisecond-scale evolution of <u>electrical pulses</u> in tens of thousands of neurons in a cubic millimeter of brain tissue.

"The thing that is novel about this work is the <u>recording density</u>," said Xie, an associate professor of electrical and computer engineering at Rice and a core member of the Rice Neuroengineering Initiative. "Microcircuits in the brain are very mysterious. We don't have many ways to map their activity, especially volumetrically. We want to deliver very dense recordings of the cortex because those are important, scientifically, for understanding how brain circuits work."

Xie collaborated on the study with colleagues from Rice and the University of California, San Francisco, including Loren Frank of UCSF and co-corresponding author Lan Luan of Rice.

Neurons are small. Each cubic millimeter of brain tissue contains about 100,000. That density is roughly the same for humans and other mammals, including the rodents that are the subject of experiments in Xie's lab. The processing power of the brain arises from synaptic connections between neurons. Synaptically linked neuron pairs are connected by narrow bridges of tissue called axons, which are just a few millionths of a meter in diameter.



Xie's team has spent years developing a material called nanoelectronic thread (NET) that is thin, ultraflexible and biocompatible, a trifecta of properties for making minimally invasive electrode implants. In previous studies, Xie's team has demonstrated techniques for emplanting tightly packed NET arrays of up to 128 electrodes. The researchers also showed their arrays could stay in place for up to 10 months, recording the pulsed spikes of electricity, or action potentials, in nearby neurons.

"When neurons fire <u>action potentials</u>, there are very faint electrical signals coming out of them," Xie said. "You have to place the electrodes very close to each neuron in order to capture that signal. Usually, that means a distance less than 100 microns."





A micro-CT scan shows some of the 1,024 flexible electrodes in a 3D neural array implanted in the visual cortex of a rodent. Credit: Xie Lab/Rice University

Using electrodes to record neuronal spikes has been a primary technique in neuroscience for decades, but the evolution of electrode materials has gradually transformed the implantation of neural electrodes from highly invasive procedures that damaged the very brain tissue the electrodes were meant to measure to procedures that result in no measurable tissue damage.



One of the primary focuses of Xie's lab is scaling up the size of its implant arrays. In the new study, Xie and colleagues, including Hanlin Zhu, one of the lead graduate students on the project, implanted arrays of 1,024 NET electrodes in a 1 cubic millimeter volume of brain tissue.

"The primary signals we try to measure are the electrical spikes coming from neurons," Xie said. "That's how they communicate. And one thing we care a lot about and really want to understand is how neurons are connected."

Xie said there is no straightforward way to probe synaptic connections.

"Axons can be very long, and each neuron can be connected by many thousands of others," he said. "It's a very, very, very messy network. And probing it is an extremely challenging task, especially while the brain is working."

The density of the new electrode array, together with its ability to capture millisecond-by-millisecond changes in the electrical spikes of individual neurons allowed Xie and co-authors to decipher potential synaptic links between neuron pairs.

"When the synapse works, you usually see a typical pattern when you look at the firing activity of the two neurons," Xie said.

It takes a bit of time for the electrical impulse that starts in the presynaptic neurons to propagate down the axon and activate the postsynaptic neuron, he said.

"We record many, many spikes, and then we need to sort the spikes and attribute each of them to individual neurons," he said. "We know the location of each electrode, or channel. And each channel records no more than a few neurons at a time. Each neuron is also typically



recorded by more than one contact as well. So, you can do something akin to triangulation to identify the location of individual neurons."

Once the neurons are mapped, it's relatively easy to calculate the distance between them and from that, the propagation time for synaptic activation.

The 1,024-electrode array gave Xie's team a ratio of approximately one electrode per 100 neurons in the cubic millimeter volume of brain tissue under study. The lab is working to create denser arrays that pack more electrodes into the same volume.

The vast majority of neurons in people's brains are unused, despite the fact that our brains typically consume about as much energy as can be supplied by the body. Neuroscientists don't fully understand why the brain has so many unused neurons, and Xie said that's a factor his team considers in the design of their <u>electrode</u> arrays.

"I want to capture as much of the interactivity as possible," he said. "I would argue that we don't need a 1-to-1 ratio of electrodes to <u>neurons</u> to capture all of it, and it is indeed my dream to capture all the interactivity."

**More information:** Zhengtuo Zhao et al, Ultraflexible electrode arrays for months-long high-density electrophysiological mapping of thousands of neurons in rodents, *Nature Biomedical Engineering* (2022). DOI: 10.1038/s41551-022-00941-y

Provided by Rice University

Citation: Engineers record neurons to pinpoint synaptic links (2022, October 19) retrieved 3 May



2024 from https://medicalxpress.com/news/2022-10-neurons-synaptic-links.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.