


Review shows neurostimulation is a promising dysphagia treatment

October 10 2022, by Mike Addelman



Summary of cerebral regions found to be activated during swallowing. The percentage represents the frequency of occurrence across 30 functional neuroimaging studies. Credit: *Journal of the Neurological Sciences* (2022). DOI: 10.1016/j.jns.2022.120434

Three types of neurostimulation technique could have the potential to help people who have the difficulty in swallowing caused by stroke or other neurological diseases, a review of 174 animal and human studies

has shown.

However, Dr. Ivy Cheng, a University of Manchester research associate who had reviewed evidence from more than 30 randomized controlled trials, says there is unfortunately only limited evidence to support the efficacy of traditional swallowing therapy used by speech therapists for [dysphagia](#).

One such neurostimulation approach is pharyngeal electrical stimulation (PES). This involves the insertion of electrodes through the nose or mouth which delivers a constant, low current to the pharynx—an important muscle in the execution of swallowing. The resultant stimulation has been shown to increase activity in the brain areas controlling swallowing.

By comparison, [repetitive transcranial magnetic stimulation](#) (rTMS) applies pulsing magnetic fields via a coil that is placed on the scalp to directly excite the underlying [brain areas](#) involved in swallowing.

A third treatment is [transcranial direct current stimulation](#) (tDCS) which uses a direct electric current to stimulate the brain directly, but this time using pads placed over the scalp in a certain orientation which, like a battery, has both anodal and cathodal configurations.

Her review found all three experimental treatments showed therapeutic potential for patients with dysphagia caused by stroke and other neurological causes and were tolerable without serious adverse effects.

However, the review—published in the *Journal of the Neurological Sciences*—shows the variability in responsiveness to the treatment has hindered its translation into [clinical practice](#) and there is uncertainty as to how long these treatment effects might last in the medium to longer term.

"Dysphagia is a condition which makes the everyday act of eating or drinking tortuous for patients," said Dr. Cheng.

"This review shows neurostimulation has a great deal of potential, but we need long term studies to assess the best protocols for using neurostimulation in treating dysphagia.

"Which is why the news of a large clinical trial of pharyngeal electrical stimulation for post-stroke dysphagia, which is running from 2021 to 2025 involving 50 sites over 4 countries, including the U.K., is so exciting."

Led by The University of Nottingham, some of the work will be carried out by Dr. Cheng's colleague Professor Shaheen Hamdy from The University of Manchester.

Dysphagia occurs when some of the 50 muscles involved in swallowing do not work to close off the windpipe to protect the lungs so food and fluid trickle through vocal cords into the lungs. Patients suffer from coughing and choking, as well as pain and voice problems when eating and drinking. Many avoid eating and others have recurrent chest infections.

Current treatment include using food thickeners, liquidizing meals and referral to speech and language therapist (SLT) for swallowing therapy. In the past, scientists thought that dysphagia was caused by problems in the [brain stem](#) alone, however more recent work has shown it is also linked to problems in the cerebral cortex.

When somebody suffers damage to the brain caused by stroke, Parkinson's, or other conditions, the part which controls swallowing might be damaged. However the evidence reviewed by Dr. Cheng showed that other parts of the brain can grow to compensate for the

damaged areas– a phenomenon called neuroplasticity.

Thanks to neuroplasticity, neural pathways in the brain get stronger if they are used regularly. If a pathway is not used regularly, it disappears.

Neurostimulation, the studies showed, can help to promote this reorganization.

More information: Ivy Cheng et al, Cerebral control of swallowing: An update on neurobehavioral evidence, *Journal of the Neurological Sciences* (2022). [DOI: 10.1016/j.jns.2022.120434](https://doi.org/10.1016/j.jns.2022.120434)

Provided by University of Manchester

Citation: Review shows neurostimulation is a promising dysphagia treatment (2022, October 10) retrieved 5 May 2024 from <https://medicalxpress.com/news/2022-10-neurostimulation-dysphagia-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--