

# Study proposes new parameters for diagnosis of sarcopenia

October 20 2022, by Maria Fernanda Ziegler

---



Measuring handgrip strength is one of the main ways of detecting sarcopenia, a syndrome characterized by loss of muscle mass, force and function. Researchers in Brazil and the UK recommended raising the minimum value considered normal, after analyzing data for more than 6,000 men and women aged 60 and over. Credit: Pixabay

Sarcopenia, a clinical syndrome characterized by progressive and extensive decline in skeletal muscle mass, force and function, is widely

considered part of aging. Early diagnosis is extremely important and begins with handgrip measurement using a dynamometer.

A recent study by researchers at the Federal University of São Carlos (UFSCar) in São Paulo state, Brazil, collaborating with colleagues at University College London (UCL) in the United Kingdom, concluded that the diagnosis protocol should be changed by raising the handgrip strength cutoff point used to detect [muscle weakness](#). They say new criteria proposed in their paper would be better predictors of [mortality risk](#) in [older adults](#), enabling healthcare professionals to detect the onset of [sarcopenia](#) earlier and more accurately.

The researchers compared cutoff points proposed in previous studies on the subject. Their analysis was based on data from the English Longitudinal Study of Ageing (ELSA) relating to 6,182 men and women aged 60 or more who were tracked for 14 years.

The UFSCar/UCL study was reported in an article published in the journal *Age and Ageing*.

The researchers took as their benchmark the diagnostic definitions issued in 2010 and revised in 2019 by the European Working Group on Sarcopenia in Older People (EWGSOP), according to which the cutoff point for handgrip strength is 27 kg for men and 16 kg for women. The article recommends raising it to 36 kg and 23 kg respectively.

"We found that lower cutoffs aren't acceptable as predictors of mortality risk. The purpose of the new reference values is to detect the risk of death as early as possible. When it's detected late, interventions such as prescribing dietary changes and resistance exercises are much harder to do. For this reason, it's important for our suggestion to be accepted by the scientific community and become a new consensus for diagnosis of sarcopenia," said Tiago da Silva Alexandre, last author of the article.

Alexandre is a professor in UFSCar's Department of Gerontology and a visiting researcher at UCL.

The lower cutoff value recommended in 2019 has always been questioned by the academic community. "Other studies had demonstrated that there were problems with the approach to diagnosing sarcopenia, but we were the first to present cutoffs capable of predicting mortality risk and compare them with those in use," he said.

By means of this comparison, the researchers were able to understand why the 2019 criteria were not working well. "Besides failing to predict mortality risk, the way sarcopenia is diagnosed underestimates the prevalence of the condition. We showed that when the cutoffs were lower, gait speed was a better metric for assessing the risk of death. However, this criterion captures only cases of advanced sarcopenia, for which prevention may no longer be possible," said Maria Claudia Bernardes Spexoto, first author of the article and a professor at the Federal University of Greater Dourados's School of Health Sciences (FCS-UFGD) in Mato Grosso do Sul state, Brazil.

Loss of muscle mass, force and function has wider implications than problems with locomotion, lack of autonomy and the risk of falling in older people. "Muscles are associated with many more functions than contracting and producing force. They participate in the immune and endocrine processes. For example, they help control certain infections via the immune system. They also help control blood sugar," Alexandre said.

Muscle quality, as well as mass, should therefore be monitored. "The skeletal muscle system participates in several other systems in the organism. If older people have good muscle mass and quality, they will have a good old age. There's no doubt about it," he said.

## **Risk of death**

According to the results of the new study, the risk of death for older people with "probable sarcopenia," or pre-sarcopenia, defined by EWGSOP as reduced muscle strength with normal muscle mass, was 30% higher than for non-sarcopenic subjects. Mortality risk was 48% higher for those with sarcopenia and 78% higher for those with severe sarcopenia.

It is worth noting that the study considered all-cause mortality and the results were adjusted for covariates (independent variables) such as sex, age, household income, marital status, schooling, smoking, alcohol intake, physical activity, arterial hypertension, diabetes, cancer, lung disease, heart disease, stroke, falls, depressive symptoms, memory status, number of medications and abdominal obesity.

According to Alexandre, the difficulty of predicting mortality risk from lower cutoffs is due to the monitoring period in the studies. While previous surveys tracked seniors for at most six years, the follow-up period used by the UFSCar/UCL team was 14 years.

"Studies proposing lower cutoffs either didn't estimate mortality risk or used data for very short periods of monitoring, so that the results were skewed," Alexandre said. "Obviously, individuals with lower muscle strength die faster, preventing those at risk in the medium to long term from being identified with precision."

Loss of muscle mass, force and function does not occur overnight but gradually over a period of years. "The natural history of sarcopenia is that it happens slowly, one day after another," he said.

## **Prevalence underestimated**

The impact of the new cutoffs proposed by the UFSCar/UCL team is significant. Among other things, it will affect sarcopenia prevalence statistics. "Higher cutoffs mean higher prevalence and more patients having to be screened so that the condition is detected well before it becomes a mortality risk," Alexandre said.

The differences in the EWGSOP guidelines set in 2010 and 2019 have led to discrepancies in prevalence, which ranges from 11% to 27.7% when the former are used and from 4.6% to 13.6% when the latter are applied.

The researchers calculated higher prevalence values based on cutoffs of 36 kg and 23 kg: 33.9% for pre-sarcopenia, 6.2% for confirmed sarcopenia and 8.6% for severe sarcopenia. "These are high rates of prevalence for a condition that can be managed so as to improve older people's quality of life and avoid premature death if it's diagnosed and treated opportunely," Alexandre said. "We're now planning a new study based on these cutoffs to estimate the prevalence of sarcopenia in Brazilians over 60."

## **New metric**

The *Age and Ageing* article recommends measuring gait speed as well as handgrip strength for early diagnosis of sarcopenia. Slowness of gait can predict a risk of frailty in older people.

"However, gait speed is itself an outcome. Older people only walk very slowly [0.8 m per second or less] when they're already weak. The new cutoffs for diagnosing sarcopenia give us an early predictor of mortality risk, which is essential to clinical practice," Alexandre said.

**More information:** Maria Claudia Bernardes Spexoto et al, European Working Group on Sarcopenia in Older People 2010 (EWGSOP1) and

2019 (EWGSOP2) criteria or slowness: which is the best predictor of mortality risk in older adults?, *Age and Ageing* (2022). DOI: [10.1093/ageing/afac164](https://doi.org/10.1093/ageing/afac164)

Provided by FAPESP

Citation: Study proposes new parameters for diagnosis of sarcopenia (2022, October 20) retrieved 24 March 2023 from <https://medicalxpress.com/news/2022-10-parameters-diagnosis-sarcopenia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.