


# Study reveals disparities in prevalence of multiple chronic conditions

October 3 2022, by Cesar Caraballo-Cordovez and Harlan Krumholz



Trends in adjusted multimorbidity prevalence by race and ethnicity, 1999–2018. Data source is the National Health Interview Survey from 1999 to 2018. Multimorbidity was defined as the presence of 2 or more of the following conditions: asthma, cancer, chronic obstructive pulmonary disease, diabetes, heart disease, hypertension, stroke, liver disease, and “weak or failing kidneys.” Annual prevalence for each race and ethnicity was obtained using logistic regression models adjusted by age, sex, and US region. The 95% confidence intervals are represented in brackets. Credit: *The American Journal of Medicine* (2022). DOI: 10.1016/j.amjmed.2022.04.010

Using data from the National Health Interview Survey, a Yale study found persistent racial and ethnic disparities in multimorbidity over a 20-year period.

A Yale study found persistent racial and [ethnic disparities](#) in multimorbidity, which is defined as the presence of two or more concurrent chronic health conditions, in the United States over a 20-year period.

Using data from the National Health Interview Survey from 1999 to 2018, investigators from the Yale Center for Outcomes Research and Evaluation (CORE) found that the annual prevalence of important chronic conditions were persistently higher among Black Americans compared with Asian, Latino/Hispanic, and [white people](#).

The study was published Sept. 12 in *The American Journal of Medicine*.

"These conditions pose a health challenge for people," said Dr. Cesar Caraballo-Cordovez, a postdoctoral associate at Yale School of Medicine and the lead author of the study. "And we found that the presence of many such conditions is persistently more common among Black people. In addition, those with low income also had the highest prevalence of these chronic conditions, overall and within the same race and ethnic group."

Also, researchers found that these chronic conditions developed earlier in life among Black individuals. "Young and middle-aged Black adults had a burden of chronic conditions equivalent to people from other racial and ethnic groups aged 5–10 years older," said Caraballo-Cordovez.

"Preventing the development of these [chronic conditions](#) should be a public health priority, and these findings reveal that [social factors](#) such

as race and income are associated with chronic disease burdens," said senior author Dr. Harlan M. Krumholz, the Harold H. Hines Jr. Professor of Medicine at Yale and director of CORE. "The findings indicate that unless we address the susceptibility of these populations to the early onset of these chronic condition, we are unlikely to address the health disparities that are so prevalent in our society."

**More information:** César Caraballo et al, Temporal Trends in Racial and Ethnic Disparities in Multimorbidity Prevalence in the United States, 1999-2018, *The American Journal of Medicine* (2022). [DOI: 10.1016/j.amjmed.2022.04.010](https://doi.org/10.1016/j.amjmed.2022.04.010)

Provided by Yale University

Citation: Study reveals disparities in prevalence of multiple chronic conditions (2022, October 3) retrieved 19 April 2024 from <https://medicalxpress.com/news/2022-10-reveals-disparities-prevalence-multiple-chronic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.