

Traumatic experiences may be associated with lower levels of sex hormones

October 12 2022



Credit: Unsplash/CC0 Public Domain

Traumatic experiences are associated with a number of adverse mental and physical health outcomes. A new study suggests that they may also be associated with lower concentrations of sex hormones in midlife

women—especially women with shorter sleep. Study results will be presented during The North American Menopause Society (NAMS) Annual Meeting in Atlanta, October 12-15, 2022.

Previous research has shown that [psychological trauma](#) has the potential to suppress ovarian function and reduce ovarian [estrogen](#) secretion. However, the relationship between [trauma](#) and sex hormones in midlife women remains largely unknown. A new study from the University of Pittsburgh involving 260 [postmenopausal women](#) has evaluated whether traumatic experiences are associated with levels of estrogens (estradiol, estrone) and follicle stimulating [hormone](#) (FSH) and whether this association is affected by sleep duration.

The researchers found that women with a history of trauma had lower levels of estrogens, including estradiol and estrone, compared with women without such history. There was no relationship between trauma and FSH levels. Findings were not accounted for by depressive or posttraumatic stress symptoms, vasomotor symptoms, or how long a woman had been postmenopausal. The relationship between trauma and hormones depended on how much women were sleeping: women with a history of trauma who were sleeping fewer than 6 hours per night had particularly low levels of estrogens.

"This work highlights the importance of trauma in relation to health at midlife, particularly given the sensitivity of women's health to hormones," says Dr. Mary Carson, lead author of the study, from the University of Pittsburgh.

"This study demonstrates the need for health care professionals to have a good understanding of a patient's history, including any traumatic experiences. This history could help identify women at increased risk for certain health issues and allow for adoption of preventive strategies," says Dr. Stephanie Faubion, NAMS medical director.

Provided by The North American Menopause Society

Citation: Traumatic experiences may be associated with lower levels of sex hormones (2022, October 12) retrieved 5 May 2024 from <https://medicalxpress.com/news/2022-10-traumatic-sex-hormones.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.